TRAUMA=NOT ONLY FRACTURES

ORTHOPEDIC=NON TRAUMATIC LIMB

-A fracture is a break in structural continuity of bone.

-If the overlying skin remains intact it is ***closed*** *(or simple).*

-if the skin or one of the body cavities is breached it is an ***open*** *(or compound) fracture*

Fractures result from:

(1) injury (single impact) (direct or indirect)

(2) repetitive stress(repetitive microtrauma)

(3) abnormal weakening of the bone (pathological Fracture)



large area
of bone
destruction
fracture

Direct injury=transverse or comminuted .
Indirect , trauma away from the fracture





Stress fracture

-Bone is dynamic both mechanical and biochemical.

-repetitive microtrauma, no time for adequate bone healing.

-athletics , dancers and military persons.



Incomplete fracture



.also called (green stick) fracture.

- fracture of cortex but still intact periostium
- occur in children due to
  - thick periostium and
  - relatively elastic bone cortex.

-usually is stable after reduction , periostium act as internal splintage.

CLASSIFICATION OF FRACTURES

1/ open or closed fracture.
2/ traumatic , stress or pathological.
3/ complete or incomplete(green stick).

For each fracture there is one or more system of classification it helps for:

1/affect modality of treatment.
2/affect the prognosis.

HOW FRACTURES ARE DISPLACED

1 /Translation (shift), *either sideway shift or overlape or impaction.*

perfect reduction is not manditary for union , some bone contact is sufficient.

2/ Angulation (tilt) .

3/ Rotation (twist)





Source: Lynn N. McKinnis: *Fundamentals of Musculoskeletal Imaging*, 4th Edition:
www.FADavisPTCollection.com
Copyright © F. A. Davis Company. All rights reserved.

HOW FRACTURES HEAL

-absolute immobilization(no mobility)at fracture site is not mandatory for healing.

-some micro movement is helpful to enhance callus formation.

-modern modalities of fracture fixation adopt this philosophy.(dynamic fixation).

-splintage to ensure
union in good position ,
alleviate pain
and permit early movement.


1. Tissue destruction & hematoma formation.
2. Inflammation & cellular proliferation.
3. Callus formation (soft callus).
4. Consolidation (woven bone).
5. Remodeling (lamellar bone).

HEALING BY CALLUS

1. *Tissue destruction and* ***haematoma*** *formation , bone die 1-2 mm from fracture line.*

2. ***Inflammation*** *and cellular proliferation – Within 8* hours of the fracture there is an acute inflammatory reaction with migration of inflammatory cells.

-The clotted haematoma is slowly absorbed and fine new capillaries grow into the area.

STAGES OF BONE FRACTURE REPAIR



3. *Callus formation* – *The differentiating stem cells* provide chrondrogenic and osteogenic. The thick cellular mass, with its islands of immature bone and cartilage, forms the callus cell population

-As the immature fiber bone (or 'woven' bone) becomes

more densely mineralized,

movement at the fracture site decreases progressively.

4. *Consolidation* – *With continuing osteoclastic and* osteoblastic activity the woven bone is transformed into lamellar bone.
-slow process and it may be several months before the bone is strong enough to carry normal loads.

5. Remodelling

Over a period of months, or even years, this crude 'weld' is reshaped by a continuous process of alternating bone resorption and formation

Types of fracture healing.

1/Gap healing

- callus formed in response to some movement, the fracture site undergo gradual load.

2/Contact healing ,In very rigid immobilization ,no callus is formed and the union will occur between fracture ends.

- e.g in fractures fixed by metal plates

UNION, CONSOLIDATION AND NON-UNION


Consolidation
-a complete repair.
-the calcified callus is ossified.
-Clinically the fracture site is not tender.
-no movement can be obtained
- attempted angulation is painless
-X-rays show the fracture line to be almost obliterated
crossed by bone trabeculae , with well-defined callus
around it.
-Repair is complete and further protection is unnecessary

Union –

-*Union is incomplete repair.*

-*the ensheathing* callus is calcified.

-Clinically the fracture site is still a little tender

-the bone moves in one piece

-attempted angulation is painful.

-X-Rays show the fracture line still clearly visible, with fluffy callus around it.

-Repair is incomplete and it is not safe to subject the unprotected bone to stress.

– Consolidation

-*is complete repair.*

-the calcified callus is ossified.

-Clinically the fracture site is not tender.

-no movement can be obtained

- attempted angulation is painless.

-X-rays show the fracture line to be almost obliterated crossed by bone trabeculae , with well-defined callus around it.

-Repair is complete and further protection is unnecessary.

Fracture repair (a) Fracture; (b) union; (c) consolidation; (d) bone remodelling. The fracture must be protected until consolidated.

non-union





Causes are:
(1) distraction and separation of the fragments, sometimes the result of interposition of soft tissues between the fragments.
(2) excessive movement at the fracture line.
(3) a severe injury that renders the local tissues nonviable or nearly so.
(4) a poor local blood supply
(5) infection.
(6) surgical intervention, if ill-judged, is another cause.

Non-unions

-septic or aseptic.
- In the aseptic, they can be either stiff or mobile as judged by clinical examination. The mobile ones can be as free and painless as to give the impression of a false joint (*pseudoarthrosis*).
- *On x-ray, non-unions are typified by* a lucent line still present between the bone fragments; sometimes there is exuberant callus trying – but failing to bridge the gap (*hypertrophic non-union*) *or at* times none at all (*atrophic non-union*).

25.7 Non-union (a,b) This patient has an obvious pseudarthrosis of the humerus. The x-ray shows a typical hypertrophic non-union. (c) Non-union associated with implant fatigue failure.

Expected time for fracture healing

-spiral+child+upper limb= three weeks.
-multiply x 2 for transverse fracture.
-multiply by another 2 for consolidation.
- multiply by another 2 for lower limb
- multiply by another 2 for adults

CLINICAL FEATURES

HISTORY:

-The fracture is not always at the site of the injury.

-If a fracture occurs with trivial trauma suspect a pathological lesion.

-Pain, bruising and swelling occur in fracture and sprains.

-*Deformity is much more suggestive*.

Clinical examination

A/ Proper assessment of ABC.

B/Exclude more serious injuries start with following sequence:

- chest injury.
- abdomen and pelvis.
- head and spine.
- limb injury.

C/ limb examination

- **-with following priorities:**
- 1- vascular examination for acute insufficiency.
- 2-examination for soft tissue injury (Gustilo and tscherene classification).
- 3-bone and joint examinaion (look , feel but try **not** to move).
- 4-tendon and nerve examination.


Gustillo classification: -
Type I - wound ≤ 1 cm.
Type II - wound 1-10cm.
Type III A: - wound > 10 cm, high energy, adequate tissue for coverage includes segmental / comminuted fractures even if wound <10cm (farm injuries are automatically Gustillo III).
Type III B: - Extensive periosteal stripping and requires free soft tissue transfer.
Type III C: - Vascular injury requiring vascular repair.



Look , feel and move

Look: skin wound,color,swelling and deformity and posture of the limb.

Feel:tenerness,crepitus,pulse and capillary filling.

Move: try to avoid unnecessary movement.

X-RAY

Remember the *rule of two,s:*

1- *Two views* at least AP and lateral.

2- *Two joints*(proximal and distal).

3- *Two limbs* , to compare in suspicion. In children, the appearance of immature epiphysis may be confused the diagnosis of a fracture

2- *Two injuries* , suspected in severe injury.

3 - *Two occasions* , some fracture appears after two weeks like fracture scaphoid.

scaphoid, lateral malleolus & femoral neck fractures.

(Painful base of thumb | tender anatomic snuff-box | pronation followed by ulnar deviation produces pain).



23.5 X-ray examination – the rule of twos. X-ray examination must be adequate. (a,b) *Two films* of the same tibia: the fracture may be 'invisible' in one view and perfectly plain in another at right angles to that. (c,d) *Two occasions*: a fractured scaphoid may not be obvious on the day of injury, but clearly seen 2 weeks later. (e,f) *Two joints*: the first x-ray (e) did not include the elbow. This was, in fact, a Monteggia fracture – the head of the radius is dislocated (f). (g,h) *Two limbs*: often the abnormality can be appreciated only by comparison with the normal side; in this case there is a fracture of the lateral condyle of the left humerus (h).

DESCRIPTION of fractures

(1) Is it open or closed?
(2) Which bone is broken , and where?
(3) Has it involved a joint surface?
(4) What is the shape of the break?(pattern)
(5) Is it stable or unstable?
(6) Is it a high-energy or a low-energy

pattern of the fracture

A transverse fracture *:is slow to join because the area* of contact is small. the fracture is stable on compression.

A spiral fracture *joins more rapidly (because* the contact area is large) but is not stable on compression.

Comminuted fractures *are often slow to join* because:
(1) they are associated with more severe soft tissue damage
(2) they are likely to be unstable.

Displacement

1. *Shift or translation* – *backwards, forwards,* sideways, or longitudinally with impaction or overlap.
2. *Tilt or angulations* – *sideways, backwards or* forwards.
3. *Twist or rotation* – *in any direction.*

Tscherne (Oestern and Tscherne, 1984)

-**classification of closed injuries:**

- *Grade 0* – *a simple fracture with little or no soft tissue* injury.
- *Grade 1* – *a fracture with superficial abrasion or* bruising of the skin and subcutaneous tissue.
- *Grade 2* – *a more severe fracture with deep soft tissue* contusion and swelling.
- *Grade 3* – *a severe injury with marked soft-tissue* damage and a threatened compartment syndrome.

Gustilo's classification of open fractures

Type I *– The wound is usually a small, clean puncture* through which a bone spike has protruded. There is little soft-tissue damage with no crushing and the fracture is not comminuted (i.e. a low-energy fracture).

Type II – *The wound is more than 1 cm long, but* there is no skin flap. There is not much soft-tissue damage and no more than moderate crushing or comminution of the fracture (also a low- to moderate-energy fracture).

Type III –

-*There is a large laceration, extensive* damage to skin and underlying soft tissue

-in the most severe examples, vascular compromise.

-the injury is caused by high-energy transfer to the bone and soft tissues.

-Contamination can be significant.

There are three grades of severity:
Type III A the fractured bone can be adequately covered by soft tissue despite the laceration.

Type III B there is extensive periosteal stripping and fracture cover is not possible without use of local or distant flaps.

-Type III C if there is an arterial injury that needs to be repaired, regardless of the amount of other soft-tissue damage

TABLE 1. Open Fractures—Gustilo Classification[1,2]

Type I	Open fracture with a skin wound <1 cm in length and clean.
Type II	Open fracture with a laceration >1 cm in length without extensive soft tissue damage, flaps, or avulsions.
Type III	Open segmental fracture with >10 cm wound with extensive soft tissue injury or a traumatic amputation (special categories in Type III include gunshot fractures and open fractures caused by farm injuries).
III_A	Adequate soft tissue coverage.
III_B	Significant soft tissue loss with exposed bone that requires soft tissue transfer to achieve coverage.
III_C	Associated vascular injury that requires repair for limb preservation.

Take home message

1/Treat the patient ,not only the fracture.
2/you should know how to describe the fracture.

THANK YOU

Treatment of fractures

Treat the patient ,not only the fracture

1/ABC.

2/ MANAGE MORE SERIOUS INJURIES.

3/LIMB INJURY MANAGEMENT ACCORDING TO PRIORITIES (BONE AND JOINT IS THE THIRD PRIORITY AFTER VASCULAR AND SOFT TISSUES




C/ limb examination
-with following priorities:
1- vascular examination for acute insufficiency
2-examination for soft tissue injury(Gustilo and tscherene classification).
3-bone and joint examination (look , feel but try not to move).
4-tendon and nerve examination.

Treatment of fracture

1/early measures(adopted for all fractures)

-ice packs

-elevation to decrease the edema

-analgesics , avoid narcotics

-temporary immobilization to

decrease pain, prevent soft tissue injury and facilitate patient transfer.

2/definitive treatment (reduction + hold + exercise)

-it is either conservative or operative

- **Reduce**.(closed or open reduction)
- **Hold.**(till fracture union)
- **Exercise**.,(of the non immobilized joints, including weight bearing

HOLD VERSUS MOVE
SPEED VERSUS SAFETY



REDUCTION

reduction is unnecessary in :

(1) when there is little or no displacement.

(2) when displacement does not matter initially (e.g. in fractures of the clavicle) .



(3) when reduction is unlikely to succeed (e.g. with compression fractures of the vertebrae).

Reduction should aim for :

1/adequate apposition:

-gap may cause delayed union and non union, the greater contact area the faster union

2/normal alignment of the bone fragments:

-some types of mal alignment is acceptable in some bones(like clavicle) and in children due to remodelling, but not the rotation.





CLOSED REDUCTION

Under appropriate anaesthesia and muscle relaxation:-

three-fold manoeuvre:

(1) the distal part of the limb is pulled in the line of the Bone.(traction and counter traction)

(2) as the fragments disengage, they are repositioned (by reversing the original direction of force if this can be reduced)

(3) alignment is adjusted in each plane.







Closed reduction is used for:

1/ all minimally displaced fractures.

2/ for most fractures in children

3/ for fractures that are stable after reduction and can be held in some form of splint or cast.

4/Unstable fractures can also be reduced using closed methods prior to stabilization with internal or external fixation. (eg. percut. Wiring)

This avoids direct manipulation of the fracture site by open reduction, which damages the local blood supply and may lead to slower healing times; increasingly

A

B


C

OPEN REDUCTION

(1) when closed reduction fails, either because of difficulty in controlling the fragments or because soft tissues are interposed between them.

(2) when there is a large articular fragment that needs accurate positioning .

(3) for traction (avulsion) fractures in which the fragments are held apart..

HOLD REDUCTION

The available methods of holding reduction are:

- Continuous traction.



- Cast splintage.



hemi hydrated calcium sulphate CaSO4. ½ H2 O in dry form, which becomes CaSO4 .2H2 O on wetting.

- Functional bracing.

- Internal fixation.
- External fixation.



CONTINUOUS TRACTION



-Traction is safe enough, provided it is not excessive

-The problem is *speed*: *not because the fracture unite* slowly (it does not) but because lower limb traction keeps the patient in hospital.

-as soon as the fracture is 'sticky' (deformable but not displaceable), traction should be replaced by bracing.

It is a method for ***reduction*** and ***holding*** the fracture



Continuous traction 'Speed' is the weak member of the quartet.

Types of traction

- ***1/Traction by gravity*** – *This applies only to upper* limb injuries. with a wrist sling the weight of the arm provides continuous traction to the humerus.



2/Skin traction – Skin traction will sustain a pull of no more than 4 or 5 kg. strapping or one way stretch Elastoplast is stuck to the shaved skin and held on with a bandage. The malleoli are protected , and cords or tapes are used for traction

3/Skeletal traction –

-*A stiff wire or pin is inserted* usually behind the tibial tubercle for hip, thigh and knee injuries, or through the calcaneum for tibial fractures – and cords tied to them for applying traction.



Fixed traction

-The pull is exerted against a fixed point.

-The usual method is to tie the traction cords to the distal end of a Thomas' splint and pull the leg down until the proximal padded ring of the splint abuts firmly against the pelvis.





Balanced traction

- The traction cords are guided over pulleys at the foot of the bed and loaded with weights; -- counter-traction is provided by the weight of the body when the foot of the bed is raised.





Methods of traction (a) Traction by gravity. (b,c,d) Skin traction: (b) fixed; (c) balanced; (d) Russell. (e) Skeletal traction with a splint and a knee-flexion piece.

Complications of traction

1/ ***Circulatory embarrassment*** *In children especially,* traction tapes and circular bandages may constrict the circulation;

2/ ***Nerve injury*** *In older people, leg traction may* predispose to peroneal nerve injury and cause a dropfoot; the limb should be checked repeatedly



3/ ***Pin site infection*** *Pin sites must be kept clean and* should be checked daily.

4/ ***skin laceration*** especially in lax skin in elderly people.

thromboembolism

CAST SPLINTAGE

hemi hydrated calcium sulphateCaSO4. ½ H2 O in dry form, which becomes CaSO4 .2H2 O on wetting.

POP

-Plaster of Paris is still widely used as a splint.

-It is safe enough, so long as one is alert to the danger of a tight cast and provided pressure sores are prevented.

-The speed of union is neither greater nor less than with traction, but the patient can go home sooner.

-joints encased in plaster cannot move and are liable to stiffen;

-While the swelling and haematoma resolve, adhesions may form that bind muscle fibers to each other and to the bone.

-Newer substitutes have some advantage over plaster (they are impervious to water, and also lighter and more radiolucent.)

Stiffness can be minimized by:

(1) delayed splintage that is, by using traction until movement has been regained, and only then applying plaster.

(2) starting with a conventional cast but, after a few weeks, when the limb can be handled without too much discomfort, replacing the cast by a functional brace which permits joint movement.

Functional bracing (cast bracing) Despite plaster the patient has excellent joint movement.



Casts 'Move' is the weakest member of the quartet.

Complications

1/Tight cast :

-The cast may be put on too tightly, or it may become tight if the limb swells. The patient complains of diffuse pain; only later – sometimes much later – do the signs of vascular compression appear. The limb should be elevated, but if the pain persists, the only safe course is to split the cast and ease it open:

(1) throughout its length and (2) through all the padding down to skin.

2/Pressure sores

-Even a well-fitting cast may press upon the skin over a bony prominence (the patella, heel, elbow or head of the ulna).

-The patient complains of localized pain precisely over the pressure spot.

-Such localized pain demands immediat inspection through a window in the cast

3/Skin abrasion or laceration

-*This is really a complication* of removing plasters, especially if an electric saw is used.

4/Loose cast

-*Once the swelling has subsided, the cast*

may no longer hold the fracture securely. If it is loose,the cast should be replaced.

INTERNAL FIXATION

-Bone fragments may be fixed with screws, a metal plate held by screws, a long intramedullary rod or nail (with or without locking screws), circumferential bands or a combination of these methods.

-Properly applied, internal fixation holds a fracture securely so that movement can begin at once; with early movement the 'fracture disease' (stiffness and edema) is abolished.


Fracture
K-wires

To stabilize a long bone fracture, a plate and screws outside the bone or a rod inside the bone may be used

Plate and screws

Intramedullary rod

A.D.A.M.

Figure-of-eight wire

©MMG 2008

-As far as speed is concerned, the patient can leave hospital as soon as the wound is healed, but he must remember that, even though the bone moves in one piece, the fracture is not united – it is merely held by a metal bridge and unprotected weight bearing is, for some time, unsafe

Safety..The greatest danger is sepsis

-if infection supervenes, all the manifest advantages of internal fixation(precise reduction, immediate stability and early movement) may be lost.

-The risk of infection depends upon:

(1) the patient – devitalized tissues, a dirty wound and an unfit patient are all dangerous.

(2) the surgeon – thorough training, a high degree of surgical dexterity and adequate assistance are all essential

(3) the facilities – a guaranteed aseptic routine, a full range of implants and staff familiar with their use are all important.



- Internal fixation 'Safety' is the weak member of the quartet.

Indications of internal fixation

1. Fractures that cannot be reduced except by operation.

2. Fractures that are inherently unstable and prone to re-displace after reduction (e.g. mid-shaft fractures of the forearm and some displaced ankle fractures).

- Also included are those avulsion fractures liable to be pulled apart by muscle action (e.g. transverse fracture of the patella or olecranon).

3. Fractures that unite poorly and slowly, principally fractures of the femoral neck.

4. Pathological fractures in which bone disease may prevent healing.

5. Multiple fractures where early fixation (by either internal or external fixation) reduces the risk of general complications and late multisystem organ failure

6. Fractures in patients who present nursing difficulties (paraplegics, those with multiple injuries and the very elderly).



(a)

(b)

(c)

Complications of internal fixation

1/Infection

Iatrogenic infection is now the most common cause of chronic osteomyelitis, the metal does not predispose to infection but the operation and quality of the patient's tissues do.

2/Non-union

-*If the bones have been fixed rigidly with a* gap between the ends, the fracture may fail to unite.

-This is more likely in the leg or the forearm if one bone is fractured and the other remains intact.

-Other causes of non-union are stripping of the soft tissues and damage to the blood supply in the course of operative fixation.

3/Implant failure

-Metal is subject to fatigue and can fail unless some union of the fracture has occurred.

-Stress must therefore be avoided and a patient with a broken tibia internally fixed should walk with crutches and stay away from partial weight bearing for 6 weeks or longer, until callus or other radiological sign of fracture healing is seen on x-ray.

4/Refracture

-*It is important not to remove metal* implants too soon, or the bone may refracture.

-A year is the minimum and 18 or 24 months safer; for several weeks after removal the bone is weak, and care or protection is needed.

EXTERNAL FIXATION



Indications

1. Fractures associated with severe soft-tissue damage (including open fractures) or those that are contaminated, where internal fixation is risky
2. Fractures around joints that are potentially suitable for internal fixation but the soft tissues are too swollen to allow safe surgery; here, a spanning external fixator provides stability until soft-tissue conditions improve.



3. Patients with severe multiple injuries, especially if there are bilateral femoral fractures, pelvic fractures with severe bleeding, and those with limb and associated chest or head injuries.

4. non united fractures, which can be excised and compressed; sometimes this is combined with bone lengthening to replace the excised segment.

5. Infected fractures, for which internal fixation might not be suitable.



© MAYO FOUNDATION FOR MEDICAL EDUCATION AND RESEARCH. ALL RIGHTS RESERVED.

The principle of external fixation is simple

- :
- -the bone is transfixed above and below the fracture with screws or tensioned wires and these are then connected to each other by rigid bars.
- -There are numerous types of external fixation devices; they vary in the technique of application
- -Most of them permit adjustment of length and alignment after application on the limb.

Complications

1/Damage to soft-tissue structures *Transfixing pins or* wires may injure nerves or vessels.

2/ joint stiffness pins may tether ligaments and muscles and inhibit joint movement..

3/Overdistraction *If there is no contact between the* fragments, union is unlikely.

4/Pin-track infection is the most common complication *,but can be less likely with good* operative technique.

EXERCISE

It is essential for :

1/decrease edema.

2/prevent joint stiffness

3/prevent muscle wasting.

4/promote fracture union.

5/enhance vascularity

Open fractures

Four golden rules:

- 1/ Antibiotic prophylaxis.
- 2/ Urgent wound and fracture debridement.
- 3/ Stabilization of the fracture.
- 4/ Early definitive wound cover.

1/Sterility and antibiotic cover

-co-amoxiclav or cefuroxime (or clindamycin if penicillin allergy) is given as soon as possible.

-cover the fracture with sterile dressing.

-

2/Debridement

-irrigation with saline aids for

removing foreign bodies,

decrease bacteria count and

prevent tissue dryness.

-wound excision removing any suspected dead tissues , skin , muscles , bones and nerves and tendon.

-may need wound extension, especially high energy open fractures

Fracture stabilization

1/ **internal fixation** can be used if we are sure of clean wound,usually after 5-7 days after negative wound swab for bacteria culture.

2/ otherwise and especially in Gustilo type III

it is safer to use **external fixation**

4/definitive wound cover

After being sure of clean wound:

1/tissue undermining and direct suturing.

2/skin graft

3/skin flap

4/myocutaneous flap

THANK
YOU

Complication of fractures

Presented by dr. Alaa Al-taee

Learning outcome

1/ how to classify the fracture complication.

2/to learn how to deal with the common complication.

Fracture complications are classified into:

1/The general complications of fractures (blood loss, shock, fat embolism cardio-respiratory failure etc.).

2/local complication of fractures:

-early complication(within first few weeks)

-late complications.

- or urgent, less urgent and late

Local complications of fractures

Urgent	Less urgent	Late
Local visceral injury	Fracture blisters	Delayed union
Vascular injury	Plaster sores	Malunion
Nerve injury	Pressure sores	Non-union
Compartment syndrome	Nerve entrapment	Avascular necrosis
Haemarthrosis	Myositis ossificans 	Muscle contracture
Infection	Ligament injury	Joint instability
Gas gangrene	Tendon lesions	Osteoarthritis
Clostridium perfringens	Joint stiffness	
	Algodystrophy	

head of the femur
scaphoid
unate
talus

1 VISCERAL INJURY

-Pneumothorax following rib fractures

- rupture of the bladder or urethra in pelvic fractures.

2 Vascular injury

Table 7.1 Blood loss from common fractures

Units (litres) lost		
1–3 (0.5–1.5)	2–4 (1–2.5)	3–5 (1.5–3)
Tibia	Femur	Pelvis
Ankle	Knee	Hip
Elbow	Shoulder	
Forearm	Humerus	

-artery may be cut, torn, compressed or contused.

-outward appearance may be normal, the intima may be detached and the vessel blocked by thrombus, or a segment of artery may be in spasm.

-The effects vary from transient diminution

of blood flow to profound ischemia, tissue death and peripheral gangrene.

Common vascular injuries

Injury	Vessel
First rib fracture	Subclavian
Shoulder dislocation	Axillary
Humeral supracondylar fracture	Brachial
Elbow dislocation	Brachial
Pelvic fracture	Presacral and internal iliac
Femoral supracondylar fracture	Femoral
Knee dislocation	Popliteal
Proximal tibial	Popliteal or its branches

Clinical features

6 Ps: pulselessness, pallor, paresthesia, pain, paralysis, poikilothermia

-The patient may complain of paraesthesia or numbness in the toes or the fingers.

-The injured limb is cold, pale, or slightly cyanosed.

- the pulse is weak or absent.

- X-rays will probably show one of the 'highrisk' fractures.

-If a vascular injury is suspected an angiogram should be performed immediately; if it is positive, emergency treatment must be started without further delay.



Treatment of vascular injury

- All bandages and splints should be removed including clothes and tourniquet(after checking its time) The fracture.

-avoid joint flexion.

-temporary reduction of grossly displaced fractures

-slight elevation.



The circulation is then reassessed repeatedly over the next half hour. If there is no improvement,

the vessels must be explored by operation

A cut vessel can be sutured, or a segment may be replaced by a vein graft;

if it is thrombosed , endarterectomy may restore the blood flow.

- if vascular repair is undertaken, stable fixation is a must and where it is practicable, the fracture should be fixed internally

3 NERVE INJURY

- -particularly common with fractures of the humerus or the elbow or the knee.
- -could be compression of nerve(neuropaxia) or severe traction injury(axontemesis) or complete cut (neurotemesis)

Common nerve injuries

Injury	Nerve
Shoulder dislocation	Axillary
Humeral shaft fracture	Radial
Humeral supracondylar fracture	Radial or median
Elbow medial condyle	Ulnar
Monteggia fracture–dislocation	Posterior-interosseous
Hip dislocation	Sciatic
Knee dislocation	Peroneal

INDICATIONS FOR EARLY EXPLORATION

Nerve injury associated with open fracture

Nerve injury with fractures that need internal fixation

Presence of a concomitant vascular injury

Nerve damage diagnosed after manipulation of the fracture

4 COMPARTMENT SYNDROME

- common in forearm or leg .

- can give rise to severe ischaemia , even if there is no damage to a major vessel.

-Bleeding, oedema or inflammation (infection) may increase the pressure within one of the osseofascial compartments.

- there is reduced capillary flow, which results in muscle ischaemia , further oedema, still greater pressure and yet more profound ischaemia .

– a vicious circle that ends, after 12 hours or less, in necrosis of nerve and muscle within the compartment . Nerve is capable of regeneration but muscle, once infarcted, can never recover and is replaced by inelastic fibrous tissue (*Volkmann's ischaemic contracture*).

-A similar cascade of events may be caused by swelling of a limb inside a tight plaster cast.



Compartment Syndrome
in Children
The doctor should start by removing the
dressing and checking the extremity.

Clinical features

-High-risk injuries are

1- fractures of the elbow, forearm bones, proximal third of the tibia.

2-fractures of the hand or foot.

3- crush injuries and circumferential burns.

4-operation (usually for internal fixation)

5- infection.



The classic features of ischemia are the five Ps:

- Pain
- Paraesthesia
- Pallor
- Paralysis
- Pulselessness.
- - But remember that in compartment syndrome the ischaemia occurs at the capillary level, so pulses may still be felt and the skin may not be pale.

-The earliest of the ‘classic’ features are pain (or a ‘bursting’ sensation).

-Ischaemic muscle is highly sensitive to stretch. If the limb is unduly painful , swollen or tense, the muscles(which may be tender) should be tested by stretching them.(passively move the joint will cause severe pain)

-Confirmation of the diagnosis can be made by measuring the intracompartmental pressures.

-The difference between diastolic pressure and compartment pressure – of less than 30 mmHg is an indication for immediate compartment decompression.

Normal intra compartmental pressure is 0-10mmHg.

Acute compartment syndrome occurs if:

-Absolute pressure increase to 25-30mmHg

-Differential pressure about 30mmHg below diastolic

Treatment

Remove bandages
Elevation at level of heart
O2
Faciotomy

-The threatened compartment(or compartments) must be promptly decompressed.

- Casts, bandages an dressings must b completely removed –

-merely splitting the plaster is useless –

- the limb should be nursed flat (elevating the limb causes a further decrease in end capillary pressure and aggravates the muscle ischaemia).

-‘fasciotomy’ means opening all compartments.

- The wounds should be left open and inspected 2 days later: if there is muscle necrosis, debridement can be carried out; if

the tissues are healthy, the wounds can be sutured (without tension) or skin-grafted.

-Muscle will be dead after 4–6 hours of total ischaemia

-there is no time to lose , decision may be made on clinical assessment if no facility of compartment pressure measurement -


Compartment Syndrome of the Right Leg

5 HAEMARTHROSIS

- -Fractures involving a joint may cause acute
- haemarthrosis.
- -The joint is swollen and tense and the
- patient resists any attempt at moving it.
- -The blood should be aspirated before dealing with the fracture.

6 INFECTION

< 5 hours - 7% infection rate

> 5 hours - 38% infection rate

- -Open fractures may become infected.
- - closed fractures hardly ever do unless they are opened by operation.

Apley -Post-traumatic wound infection is now the most common cause of chronic osteomylitis.

7 GAS GANGRENE

- -is produced by clostridial (especially *Clostridium welchii*).
- -*These are* anaerobic organisms that can survive and multiply only in tissues with low oxygen tension
- -site for infection, is a dirty wound with dead
- muscle that has been closed without adequate
- debridement.
- -Toxins produced by the organisms destroy the cell wall and rapidly lead to tissue necrosis.

- -Clinical features appear within 24 hours of the
- Injury.
- - intense pain and swelling around the wound and a brownish discharge may be seen.
- - gas formation is usually not very marked.
- -There is little or no pyrexia .
- - the pulse rate is increased
- -characteristic smell becomes evident
- -Rapidly the patient becomes toxaemic and may lapse into coma and death.

-It is essential to distinguish gas gangrene, which is characterized by myonecrosis, from anaerobic cellulitis, in which superficial gas formation is abundant but toxaemia usually slight.

-Failure to recognize the difference may lead to unnecessary amputation for the non-lethal cellulitis.

Treatment

-there is no effective antitoxin against *C. welchii.*

-The key to life-saving treatment is early diagnosis.

-General measures, such as fluid replacement and intravenous antibiotics, are started immediately.

-Hyperbaric oxygen has been used as a means of limiting the spread of gangrene.

-the mainstay of treatment is proper decompression of the wound and removal of all dead tissue.

- In advanced cases, amputation may be essential.

FRACTURE BLISTERS

-Two distinct blister types are seen after fractures:

1/ clear fluid-filled vesicles

2/blood-stained ones.

-Both occur during limb swelling and are due to elevation of the epidermal layer of skin from the dermis

-There is no advantage to puncturing the blisters (it may even lead to increased local infection)

PLASTER AND PRESSURE SORES

-Plaster sores occur where skin presses directly onto bone.

-They should be prevented by padding the bony points .

-While a plaster sore is developing the patient feels localized burning pain.

- A window must immediately be cut in the plaster.

-skin necrosis proceeds unnoticed.

(a) (b)

Pressure sores Pressure sores are a sign of carelessness. (a,b) Sores from poorly supervised treatment in a Thomas splint.

INJURIES OF THE PHYSIS



-In children over 10 per cent of fractures involve injury to the growth plate (or physis).

-Because the physis is a relatively weak part of the bone :

-strains that might cause ligament injuries in adults are liable to result in separation of the physis in children.

-

-The fracture usually runs transversely through the hypertrophic or the calcified layer of the growth Plate this has little effect on longitudinal growth, which takes place in the germinal and proliferating layers of the physis.

- However, if the fracture traverses the cellular 'reproductive' layers of the physis, it may result in premature ossification of the injured part and serious disturbances of bone growth.



Classification

Salter and Harris

:

- ***Type 1*** *– A transverse fracture through the hypertrophic* or calcified zone of the plate. Even if the fracture is quite alarmingly displaced, the growing zone of the physis is usually not injured and growth disturbance is uncommon.
- ***Type 2*** *– This is essentially similar to type 1, but* towards the edge the fracture deviates away from the physis and splits off a triangular metaphyseal fragment of bone (sometimes referred to as the *Thurston– Holland fragment).*

Salter-Harris Classification

REBEL REVIEWS



I	II	III	IV	V
S	**A**	**L**	**T**	**ER**
Separated Growth Plate	Above Growth Plate	Below Growth Plate	Through Growth Plate	Erasure of Growth Plate

• ***Type 3*** – *A fracture that splits the epiphysis and then*
veers off transversely to one or the other side,
through the hypertrophic layer of the physis.
Inevitably it damages the 'reproductive' layers of
the physis (as these layers are closer to the epiphysis
than the metaphysis) and may result in growth disturbance.

• ***Type 4*** – *As with type 3, the fracture splits the epiphysis,*
but it extends into the metaphysis. These fractures are liable to displacement and a consequent misfit between the separated parts of the physis, resulting in asymmetrical growth


I
II
III
IV
V

Type 5 –

A longitudinal compression injury of the physis.

There is no visible fracture but the growth plate is crushed and this may result in growth arrest.

Physeal injuries Type 1 – separation of the epiphysis – which usually occurs in infants but is also seen at puberty as a slipped femoral epiphysis. Type 2 – fracture through the physis and metaphysis – is the commonest; it occurs in older children and seldom results in abnormal growth. Type 3 – an intra-articular fracture of the epiphysis – needs accurate reduction to restore the joint surface. Type 4 – splitting of the physis and epiphysis – damages the articular surface and may also cause abnormal growth; if it is displaced it needs open reduction. Type 5 – crushing of the physis – may look benign but ends in arrested growth.

THANK YOU FOR LISTENING

© 2014 KeepCalmStudio.com

LATE COMPLICATIONS OF FRACTURES

DELAYED UNION

٦ أشهر

- **Causes:**
- *1/biological.*
- *2/biomechanical*
- *3/patient-related.*

BIOLOGICAL

- ***1/Inadequate blood supply:*** *especially in comminuted fractures due to interrupted indosteal and periosteal blood supply.*
- ***2/Severe soft tissue damage*** by:
- (1) reducing the effectiveness of muscle splintage
- (2) damaging the local blood supply
- (3) diminishing or eliminating the osteogenic input
- from mesenchymal stem cells within muscle.

- ***3/Periosteal stripping****, excessive stripping of*
- periosteum during internal fixation is an avoidable
- cause of delayed union.

BIOMECHANICAL

- ***1/Imperfect splintage :***
- *-Excessive traction (creating a* fracture gap)
- -excessive movement at the fracture site
- -In the forearm and leg a single-bone fracture may be held apart by an intact fellow bone.
- ***2/Over-rigid fixation***
- ***3/Infection***

PATIENT RELATED

- -uncooperative patient, alcoholic, dementia.
- -immune suppressed.
- -malnutrition, vitamin D deficiency

Clinical features

- -Fracture tenderness persists .
- -if the bone is subjected to stress, pain may be acute.
- -**On x-ray,** the fracture line remains visible and there is very little or incomplete callus formation or periosteal reaction.
- However, the bone ends are not sclerosed or atrophic.
- The appearances suggest that ,although the fracture has not united, it eventually will.

Treatment

- **CONSERVATIVE**
- fracture loading is an important stimulus to union and can be enhanced by:

  (1) encouraging muscular exercise

  (2) by weight bearing in the cast or brace.

OPERATIVE

- If union is delayed for more than 6 months and there is no sign of callus formation, internal fixation and bone grafting are indicated.
-
- The operation should be planned in such a way as to cause the least possible damage to the soft tissues

NON-UNION

- -In a minority of cases delayed union gradually turns into non-union – that is it becomes apparent that the fracture will never unite without intervention.
- -Movement can be elicited at the fracture site and pain diminishes; the fracture gap becomes a type of pseudoarthrosis

X-ray of nonunion

- -The fracture is clearly visible but the bone on either side of it may show either exuberant callus or atrophy.
-
- - so divided into hypertrophic and atrophic types.

- *1/In hypertrophic non-union ,the bone ends are*
- enlarged, suggesting that osteogenesis is still active
- but not quite capable of bridging the gap.
-
- 2/In *atrophic non-union, osteogenesis seems to have ceased. The* bone ends are tapered or rounded with no suggestion of new bone formation.

Causes of non union

- SAME CAUSES OF DELAYED UNION,BUT PROLONGED ENOUGH TO CAUSE NON UNION

Pesado arthrosis

(a)

(b)

(c)

(d)

Non-union

(a) This patient has an obvious pseudarthross of the humerus. The x-ray (b) shows a typical hypertrophic non-union. (c,d) Examples of atrophic non-union.

Treatment of non union

- **CONSERVATIVE**
- -Non-union is occasionally symptomless, needing no treatment or a removable splint.
- - if symptoms are present, operation is better than trial of prolonged splintage.

OPERATIVE treatment

- -With hypertrophic non-union and in the absence of deformity, rigid fixation alone (internal or external) may lead to union. ,no need for bone graft.

- -With atrophic non-union, fixation alone is not enough. Fibrous tissue in the fracture gap, as well as the hard, sclerotic bone ends is excised and bone grafts are packed around the fracture.

Non-union – treatment (a) This patient with fractures of the tibia and fibula was initially treated by internal fixation with a plate and screws. The fracture failed to heal, and developed the typical features of hypertrophic non-union. (b) After a further operation, using more rigid fixation (and no bone grafts), the fractures healed solidly. (c,d) This patient with atrophic non-union needed both internal fixation and bone grafts to stimulate bone formation and union (e).

(a)

(b)

(c)

(d)

Non-union – treatment by the Ilizarov technique

Hypertrophic non-unions can be treated by gradual distraction and realignment in an external fixator (a–d). Atrophic non-unions will need more surgery; the poor tissue is excised (e,f) and replaced through bone transport (g,h).

MALUNION

- When the fragments join in an unsatisfactory position (unacceptable angulation, rotation or shortening) the fracture is said to b malunited.
- **Causes are:**
- **1/** failure to reduce a fracture adequately.
- **2/** failure to hold reduction while healing proceeds.
- **3/**gradual collapse of comminuted or osteoporotic bone.

osteoporosis

Clinical features

- -The deformity is usually obvious.
- -sometimes the true extent of malunion is apparent only on x-ray.
- -Rotational deformity of the femur, tibia, humerus or forearm may be missed unless the limb is compared

with its opposite fellow.

Prevention:

- -X-rays are essential to check the position of the fracture while it is uniting. This is particularly important
- during the first 3 weeks, when the situation may
- change without warning.

TREATMENT OF MALUNION

- **1. In adults**, fractures should be reduced as near to the anatomical position as possible.
- -Angulation of more than 10–15 degrees in a long bone or a noticeable rotational deformity may need correction by osteotomy and fixation.
- **2. In children,** angular deformities near the bone
- ends (and especially if the deformity is in the same
- plane as that of movement of the nearby joint) will
- usually remodel with time; **rotational** deformities
- will not remodel.

- 3. In the lower limb, shortening of more than 2.0 cm is seldom acceptable to the patient and a limb length equalizing procedure may be indicated.
- 4. The patient's expectations (often prompted by cosmesis) may be quite different from the
- surgeon's; they are not to be ignored.

- 5. Early discussion with the patient, and a guided view of the x-rays, will help in deciding the need for treatment and may prevent later misunderstanding.

- 6. malalignment of more than 15 degrees in any plane may cause asymmetrical loading of the joint above or below and the late development of secondary osteoarthritis; this applies particularly to the large weightbearing joints.

AVASCULAR NECROSIS

- **Certain regions are notorious for avascular necrosis**:
- (1) the head of the femur (after fracture of the femoral neck or dislocation of the hip).
- (2) the proximal part of the scaphoid (after
- fracture through its waist).
- (3) the lunate (following dislocation) .
- (4) the body of the talus (after fracture
- of its neck).

Clinical features

- -initially there are no symptoms associated with avascular
- necrosis, but if the fracture fails to unite or if the bone
- collapses the patient may complain of pain.
- -X-ray shows the characteristic increase in x-ray density,
- which occurs as a consequence of two factors:
- disuse osteoporosis in the surrounding parts gives the
- impression of 'increased density' in the necrotic segment,
- and collapse of trabeculae compacts the bone
- and increases its density.

(the consequence of new bone ingrowth in the necrotic segment and disuse osteoporosis in the surrounding parts

(a) (b) (c)

Avascular necrosis (a) Displaced fractures of the femoral neck are at considerable risk of developing avascular necrosis. Despite internal fixation within a few hours of the injury (b), the head-fragment developed avascular necrosis. (c) X-ray after removal of the fixation screws.

Treatment

- -Treatment usually becomes necessary when joint
- function is threatened or the pain become severe.

- - In old people with necrosis of the femoral head an *arthroplasty* is the choice.
- -in younger people, realignment *osteotomy*(or, in some cases, *arthrodesis*) may be wiser.
- -Avascular necrosis in the scaphoid or talus may need no more than symptomatic treatment, but arthrodesis of the wrist or ankle is sometimes needed if severe pain evolved.

GROWTH DISTURBANCE

- -In children, damage to the physis may lead to abnormal or arrested growth.
- -A transverse fracture through the growth plate is not always disastrous; the fracture runs through the hypertrophic and calcified layers and not through the germinal zone, there may not be any disturbance of
- growth.
- - fractures that split the epiphysis inevitably traverse the growing portion of the physis, and so further growth may be asymmetrical and the bone end characteristically angulated.
- -if the entire physis is damaged, there may be slowing or complete cessation of growth.

BED SORES

- -Bed sores occur in elderly or paralysed patients.
- - The skin over the sacrum and heels is especially vulnerable.
- -Careful nursing and early activity can usually prevent
- bed sores; once they have developed, treatment is difficult
- – it may be necessary to excise the necrotic tissue
- and apply skin grafts.
- -In recent years vacuum-assisted closure (a form of negative pressure dressing) has been used for sacral bed sores.

Bed sores Bed sores in an elderly patient, which kept her in hospital for months.

MYOSITIS OSSIFICANS

- -Heterotopic ossification in the muscles sometimes occurs after an injury, particularly dislocation of the elbow or a blow to the brachialis, deltoid or quadriceps.
- -It is thought to be due to muscle damage, but it also occurs without a local injury in
- unconscious or paraplegic patient

Clinical features

- -Soon after the injury, the patient (usually a fit young man) complains of pain and local swelling and soft-tissue tenderness.
- -***X-ray*** is normal .
- -***bone scan*** may show increased activity.
- -Over the next 2–3 weeks the pain gradually subsides, but joint movement is limited;
- ***x-ray*** may show fluffy calcification in the soft tissues.
- -By 8 weeks the bony mass is easily palpable
- and is clearly defined in the x-ray.

Treatment

Avoid vigorous exercises
Active movements
excision
Indomethacin or radiotherapy

- -The worst treatment is to attack an injured and slightly stiff elbow with vigorous muscle-stretching exercises; this is liable to precipitate or aggravate the condition.
-
- -The joint should be rested in the position of function until pain subsides; gentle **active** movements
- are then begun.(avoid passive movement).

- -Months later, when the condition has stabilized, it
- may be helpful to excise the bony mass.

- - Indomethacin or radiotherapy should be given to help prevent a recurrence.

TENDON LESIONS

- *1/Tendinitis.*
- *2/Tendon rupture due to contineous friction with rough fracture susface*



LATE NERVE COMPRESSION

- **Common sites are:**
- (1) the ulnar nerve, due to a valgus elbow following a malunited lateral condyle or supracondylar fracture.

- (2) the median nerve, following injuries around the wrist
- (3) the posterior tibial nerve, following fractures around the ankle.

MUSCLE CONTRACTURE

- -Following arterial injury or compartment syndrome,
- the patient may develop ischaemic contractures of the
- affected muscles (*Volkmann's ischaemic contracture*).

- -the patient presents with deformity and stiffness, but numbness is inconstant.

- -The sites most commonly affected are the forearm and hand,
- leg and foot.

- -In a severe case affecting the forearm, there will be
- wasting of the forearm and hand, and clawing of the
- fingers.





JOINT INSTABILITY

- **Causes include the following:**
- • *Ligamentous laxity* – *especially at the knee, ankle* and metacarpophalangeal joint of the thumb.
- • *Muscle weakness* – *especially if splintage has been* excessive or prolonged.
- • *Bone loss* – *especially after a gunshot fracture or* severe compound injury.
- -*recurrent dislocation. The* commonest sites are:
- (1) the shoulder – if the glenoid labrum has been detached (a Bankart lesion) .
- (2) the patella – if, after traumatic dislocation, the restraining patellofemoral ligament heals poorly.

JOINT STIFFNESS

- -commonly occurs in the knee, elbow, shoulder and (worst of all) small joints of the hand.
- 1/ joint itself has been injured.
- 2/ a haemarthrosis forms and leads to synovial adhesions.
- 3/ More often the stiffness is due to oedema and fibrosis of the capsule, ligaments and muscles around the joint, or adhesions of the soft tissues to each other or to the underlying bone.

ligament => Stretch => كلش زين صحيح

- -All these conditions are made worse by prolonged immobilization; especially if the joint has been held in a position where the ligaments are at their shortest.
- - The best treatment is prevention – by exercises that keep the joints mobile from the outset.
- - If a joint has to be splinted, make sure that it is held in the 'position of safety,(max.ligament length)'

COMPLEX REGIONAL PAIN SYNDROME (ALGODYSTROPHY)

aberrant inflammatory response that leads to sustained sympathetic activity in a perpetuated reflex arc

- - *Sudeck's atrophy* is a common type of the syndrome.
- - The patient complains of continuous pain, often
- described as 'burning' in character.
- -***At first*** there is local swelling, redness and warmth, as well as tenderness and moderate stiffness of the nearby joints.
- -***As the weeks go*** by the skin becomes pale and atrophic.
- -movements are increasingly restricted and the patient
- may develop fixed deformities.
- -X-rays characteristically show patchy rarefaction of the bone

(a)

(b)

(c)

Complex regional pain syndrome (a) Regional osteoporosis is common after fractures of the extremities. The radiolucent bands seen here are typical. (b) In algodystrophy the picture is exaggerated and the soft tissues are also involved: here the right foot is somewhat swollen and the skin has become dusky, smooth and shiny. (c) In the full-blown case, x-rays show a typical patchy osteoporosis. (d) Similar changes may occur in the wrist and hand. They are always accompanied by (e) increased activity in the radionuclide scan.

- -The earlier the condition is recognized and treatment
- begun, the better the prognosis.
- -Elevation and active exercises are essential.
- -In the early stage of the condition, anti-inflammatory drugs and adequate analgesia are helpful.
- -desensitization methods, regional anaesthesia, and use of drugs like amitriptyline, carbamazepine and gabapentin may help.
- - prolonged and dedicated physiotherapy, is the
- mainstay of treatment.

OSTEOARTHRITIS

- -joint articular damage by:
- 1/injury of the hyaline cartilage during the trauma.
- 2/malalignment cause abnormal load transfer

THANK
YOU

Upper Limb T

Presented By

Dr. Karrar Al- Jubory

Orthopedic surgeon

FICMS, CABMS, (Ortho



Bony anatomy







blood supply







Nerve supply





Movement


Flexion
Extension
Medial rotation
Lateral rotation
Adduction
Abduction
Abduction
Flexion
Extension
Circumduction
Adduction

Topics

- Clavicle
- Humerus
- Forearm
- Distal Radius
- Shoulder Dislocation
- Elbow dislocation

Clavicle Fractures

- Allman Classification of Clavicle Fractures

– Type I Middle Third (80%)

– Type II Distal Third (15%)

- Differentiate whether ligaments attached to lateral or medial fragment

– Type III Medial Third (5%)

Clavicle Fractures

Mechanism

– Fall onto shoulder (87%)

– Direct blow (7%)

– Fall onto outstretched hand (6%) FOOSH

Trimodal distribution

Group I (13yrs)

Group 2 (47yrs)

Group 3 (59yrs)

The clavicle is the last ossification center to complete (sternal end) at about 22-25ys

Clinical Evaluation

– Inspect and palpate for deformity abnormal motion

– Thorough distal neurovascular exam

– Auscultate the chest for the possibility of lung injury or pneumothorax

Radiographic Exam

– AP chest radiographs.

– Clavicular 45deg A/P oblique X-rays

X-ray





Treatment

1-Less than 2cm shortening and displacement
2-Less than 1cm displacement of the superior shoulder suspensory complex
3-closed
4-no neurovascular injury
5-low demand patient

- Closed Treatment – Sling immobilization for usually ~~3-4 weeks~~ with early ROM encouraged **Rough assessment**
- Operative intervention
  - – Fractures with neurovascular injury
  - – Fractures with severe associated chest injuries
  - – Open fractures
  - – Group II, type II fractures
  - – Cosmetic reasons, uncontrolled deformity
  - – Nonunion

More than 2 cm shortening

Associated Injuries

– Brachial Plexus Injuries >> Contusions most common, penetrating (rare)

– Vascular Injury

– Rib Fractures

– Scapula Fractures

– Pneumothorax Clavicle Fractures

Shoulder Dislocations

 Epidemiology

– Anterior: Most common

– Posterior: Uncommon, 10%, Think Electrocutions & Seizures

– Inferior: Rare, hyper-abduction injury

Ant. Recurrence

– Age 20: 80-92%, 30: 60%, 40: 10-15%

Glenoid Fracture, Greater Tuberosity Fracture, Subscapularis Tear, Axillary artery injury (older pts with atherosclerosis), Axillary nerve neuropraxia

Clinical Evaluation

– Examine(deltoid function, sensation over lateral shoulder)

– Examine M/C nerve (biceps function and anterolateral forearm sensation)

 Radiographic Evaluation – True AP shoulder – Axillary Lateral – Scapular Y – Stryker Notch View (Bony Bankart)

Traumatic – Atraumatic (Congenital Laxity) – Acquired

Radiology


Normal Shoulder
(AP View)
Coracoid Process
Clavicle
Acromion Process
Humeral Head
Glenoid
Body of
Scapula
Humerus
Lateral Border
Scapula
EBM | CONSULT

Posterior Dislocation

– Adduction/Flexion/IR at time of injury
– Electrocution and Seizures cause overpull of subscapularis and latissimus dorsi

– Look for "lightbulb sign" and "vacant glenoid" sign

– Reduce with traction and gentle anterior translation

Treatment

– Nonoperative

–Traction/counter-traction

- Hippocratic technique- Effective for one person.
- Stimson technique-
- Milch Technique- Arm is abducted and externally rotated with thumb pressure applied to the humeral head
- Scapular manipulation
- Kocher methods



Stimson maneuver
Patient prone on table with affected limb hanging freely over edge; 10–15-lb weight suspended from wrist. Gradual traction overcomes muscle spasm and in most cases achieves reduction in 20–25 minutes.
Stimson Maneuver (Netter's Concise Orthopedic Anatomy 2nd Edition)

Post-reduction

– Post reduction films are a must to confirm the position of the humeral head
– Pain control

– Immobilization for 7-10 days then begin progressive ROM

 Operative Indications

– Irreducible shoulder (soft tissue interposition)

– Displaced greater tuberosity fractures

– Glenoid rim fractures bigger than 5 mm

– Elective repair for younger patients

Recurrent

Proximal Humerus Fractures

Epidemiology

– Most common fracture of the humerus

– Higher incidence in the elderly, thought to be related to osteoporosis

– Females 2:1 greater incidence than males

Mechanism of Injury

– Most commonly a fall onto an outstretched arm from standing height

– Younger patient >> Motor Vehicle Accident

Clinical Evaluation

– Patients typically present with arm held close to chest by contralateral hand.

– Pain and crepitus detected on palpation

– Careful NV exam to the axillary nerve.

Test sensation over the deltoid.

Deltoid atony does not necessarily confirm an axillary nerve injury

Radiology

Treatment

Proximal Humerus Fractures E

– Minimally displaced fractures- Sling immobilization, early motion

– Two-part fractures-

 Anatomic neck fractures likely require ORIF. High incidence of osteonecrosis

 Surgical neck fractures that are minimally displaced can be treated conservatively. Displacement usually requires ORIF

– Three-part fractures

 Due to disruption of opposing muscle forces, these are unstable so closed treatment is difficult. Displacement requires ORIF.

– Four-part fractures

 In general for displacement or unstable injuries ORIF in the young and hemiarthroplasty in the elderly and those with severe comminution. High rate of AVN (13-34%)



Humeral Shaft Fractures

- Mechanism of Injury

– Direct trauma is the most common especially MVA

– Indirect trauma such as fall on an outstretched hand

– Fracture pattern depends on stress applied
- Compressive- proximal or distal humerus
- Bending- transverse fracture of the shaft
- Torsional- spiral fracture of the shaft
- Torsion and bending- oblique fracture usually associated with a butterfly fragment

Clinical evaluation

– Thorough history and physical

– Patients typically present with pain, swelling, and deformity of the upper arm
– Careful NV exam important as the radial nerve is in close proximity to the humerus and can be injured

Treatment

Conservative

– Goal of treatment is to establish union with acceptable alignment

– >90% of humeral shaft fractures heal with nonsurgical management

- 20 degrees of anterior angulation,[1]
- 30 degrees of varus angulation[2] and up to 3[3] cm of shortening are acceptable
- Most treatment begins with application of a coaptation spint or a hanging arm cast followed by placement of a fracture brace

Operative Treatment

- Indications

Inadequate reduction, nonunion, associated injuries, open fractures, segmental fractures, associated vascular or nerve injuries

- Most commonly treated with plates and screws but also IM nails
- Holstein-Lewis Fractures – Distal 1/3 fractures – May entrap or lacerate radial nerve as the fracture passes through the intermuscular septum



Elbow Dislocations

Post

- Epidemiology 11-28% of injuries to the elbow
  - Posterior dislocations most common
  - Highest incidence in the young 10-20 years

Most commonly due to FOOSH or elbow resulting in force to unlock the olecranon from the trochlea

Posterior >> Hyperextension, valgus stress, arm abduction, and forearm supination

Anterior >> ensuing from direct force to the posterior forearm with elbow flexed

Clinical Evaluation

- Patients typically present guarding
- Usually has gross deformity and swelling
- Careful NV exam in important and should be done prior to manipulation
- Repeat after reduction



Anterior humoral line should pass central of middle third



Radiograph

Treatment

– Posterior Dislocation

- Closed reduction under sedation
- Reduction should be performed with the elbow flexed while providing distal traction
- Post reduction splint with the elbow at 90 degrees
- Open reduciton

– Anterior Dislocation

Closed reduction under sedation

humerus

- Associated injuries

Radial head fx (5-11%)

- Type I- Conservative
- Type II/III- Attempt ORIF vs. radial head replacement
- No role for solely excision of radial head



Coronoid process fractures

Medial or lateral epicondyle fx

Posterolateral rotary instability,

lateral collateral ligament (LCL) complex

Elbow radiography











Supra condylar fractures

Most common elbow fractures seen in children 5 to 6yrs

- Boys vs gals 3:2
- Left or non dominant side more commona

Mechanism

- Depending on direction of displacement of distal fragment

- Extension & flexion types
- Outstretched hand with elbow full extension

Displacements posteromedial vs posterolateral



Signs & symptoms

- History- pulling or fall
- Pain, swelling
- Tenderness-both condyles
- Decreased ROM
- Limited extension
- Gross deformity
- Anterior **pucker sign**

S-configuration





Examination

Sensory

- Radial nerve- dorsal 1st web space

  Median nerve – palmar index finger

  Ulnar nerve – palmar little finger

Motor

- Finger, wrist, thumb extension -Radial nerve

- Distal IP flexion & thumb IP flexion - Anterior interosseous nerve
- Thenar strength -Median nerve
- Interossei -Ulnar nerve

Vascular examination

- Presence of pulse, warmth, capillary refill, and colour of the hand.
- Vascular status three categories
- Hand well-perfused (warm and red), radial pulse present
- Hand well-perfused, radial pulse absent
- Hand poorly perfused (cool and blue or blanched), radial pulse absent

Radiographic evaluation

- True Ap ,Lat, Oblique views
- Initial xray may be negative except for a posterior fat pad sign



- Anterior humeral line
- Baumann angle or Humeral capitellar angle -Normal range 9 to 26 deg, - 10 degrees is acceptable

-A Increase in the Baumann angle is a sign that a fracture is in varus angulation

26.15 Supracondylar fractures X-rays showing supracondylar fractures of increasing severity. (a) Undisplaced. (b) Distal fragment posteriorly angulated but in contact. (c) Distal fragment completely separated and displaced posteriorly. (d) A rarer variety with anterior angulation.

X-rays

Undisplaced fractures are easily missed; there may be no more than subtle features of a soft-tissue haematoma. In the common *posteriorly displaced fracture* the distal fragment is tilted backwards and/or shifted backwards. In the rare *anteriorly displaced fracture* the fragment is tilted forwards.

The anteroposterior x-ray is often difficult to interpret because it is taken with the elbow flexed. The degree of sideways tilt (angulation) may therefore not be appreciated. This is where Baumann's angle is most helpful; wherever possible it should be accurately measured and compared with that of the uninjured side (Figure 26.14).

Treatment

If there is even a suspicion of a fracture, the elbow is gently splinted in 30 degrees of flexion to prevent movement and possible neurovascular injury during the x-ray examination.

Undisplaced fractures

The elbow is immobilized at 90 degrees and neutral rotation in a lightweight splint or cast and the arm is supported by a sling. *It is essential to obtain an x-ray 5–7 days later to check that there has been no displacement*. The splint is retained for 3 weeks and supervised movement is then allowed.

Posteriorly minimally angulated fractures

Swelling is usually not severe and the risk of vascular injury is low. If the posterior cortices are in continuity, the fracture can be reduced under general anaesthesia by the following stepwise manoeuvre: (1) traction for 2–3 minutes in the length of the arm with counter-traction above the elbow; (2) correction of any sideways tilt or shift and rotation (in comparison with the other arm); (3) gradual flexion of the elbow to 120 degrees, and pronation of the forearm, while maintaining traction and exerting finger pressure behind the distal fragment to correct posterior tilt. *Then feel the pulse and check the capillary return* – if the distal circulation is suspect, immediately relax the amount of elbow flexion until it improves.

X-rays are taken to confirm reduction, checking carefully to see that there is no varus or valgus angulation and no rotational deformity. If the acutely flexed position cannot be maintained without disturbing the circulation, or if the reduction is unstable, the fracture should be fixed with percutaneous crossed Kirschner wires (K-wires) (take care not to skewer the ulnar nerve!).

Following reduction, the arm is held in a collar and cuff; the circulation should be checked repeatedly during the first 24 hours. An x-ray is obtained after 3–5 days to confirm that the fracture has not slipped. If it has, do not delay – a further attempt at reduction is still possible. If reduction is satisfactory, the splint is retained for 3 weeks, after which movements are begun.

Posteriorly displaced fractures

These are usually associated with *severe swelling*, are difficult to reduce and are often unstable; moreover, there is a considerable risk of neurovascular injury or circulatory compromise due to swelling. The fracture should be reduced under general anaesthesia as soon as possible, by the method described above, and then held with percutaneous crossed K-wires; this obviates the necessity to hold the elbow acutely flexed. Care should be taken not to injure the ulnar and radial nerves. Postoperative management is the same as for simple angulated fractures.

Anteriorly displaced fractures

The fracture is reduced by pulling on the forearm with the elbow semi-flexed, applying thumb pressure over the front of the distal fragment and then extending the elbow fully. A posterior slab is bandaged on and retained for 3 weeks. Thereafter, the child is allowed to regain flexion gradually.

Treatment

TYPE I: splinted with the elbow in 90 degrees of flexion.
TYPE II: closed reduction and splinting
TYPE III: closed reduction and percutaneous pinning (K- wires)

- Initial splinting

- Duration of immobilization 3-4wks.
- • Avoid Tight bandaging, excess flexion, extension
- • Limb elevated
- • Neurologic & vascular status
- • Look for compartment syndrome

Operative

We don't use screws in pediatric

Ideally

Closed Reduction and Pinning K wire

• initially attempted - including type III fractures

• first reduced in the frontal plane

• elbow is then flexed while the olecranon is pushed anteriorly to correct the sagittal deformity

Treatment

Indications of a successful reduction

- Intact medial and lateral columns on oblique views
- Anterior humeral line passing through the middle third of the capitellum on the lateral view
- immobilized in 50 to 60 degrees of flexion

No need for any physiotheraphy

Indications for open reduction

- Gap in the fracture site
- An irreducible fracture with a rubbery feeling
- Closed reduction fails
- Fractures associated with a dysvascular limb

Treatment with Traction(conservative)

- Severe comminution • Lack of anesthesia • Medical conditions prohibiting • Malunion is common

• توضيح للنيرف انجري راح احدده من distal fragment فاذا جان post.medial راح تصير

spike of humerus في ant.laterl فيسوي انجري لل radial nerve يعني عكس اما اذا جان post.lat راح يسوي

انجري The anterior interosseous nerve a branch of the median nerve و البركيل ارتري

اذا ant displacement يصير الانجري post يعني ulnar nerve

- Vascular injury **Brachial artery**
- Compartment syndrome **Early complication Pain out of proportion**
- Neurologic defect **anterior interosseous (flexor pollicis longus, flexor digitorum profundus and pronator quadratus).**
- Elbow Stiffness **Trauma or surgery**
- Pin Track Infections
- Myositis Ossificans
- Cubitus varus “gunstock deformity”





Lateral Condyle Fracture

Lateral Condyle Fractures are the second most common fracture in the pediatric elbow and are characterized by a higher risk of nonunion, malunion, and AVN than other pediatric elbow fractures

of all distal humerus fractures 17% in the pediatric population

2nd most common elbow fracture after supracondylar

Mechanism of injury

Pull-off theory

avulsion fracture of the lateral condyle that results from the pull of the common extensor musculature

Push-off theory

fall onto an outstretched hand causes impaction of the radial head into the lateral condyle causing fracture

Milch Classification

Type I

Fracture line is lateral to trochlear groove (less common, elbow is stable as fracture does NOT enter trochlear groove)

Type II

Fracture line extends medially into trochlear groove (more common, more unstable)


Type 1
less common
stable
Salter–Harris Type IV
Trochlea
Type 2
more common
unstable
Salter–Harris Type II
Trochlea

Ossification center of the Elbow

	Years of ossification	*Years at fusion*
•Capitellum	•1	•12-14
•Radial head	•3	•14-16
•Internal (medial) epicondyle	•5	•16-18
•Trochlea	•7	•12-14
•Olecranon	•9	•15-17
•External (lateral) epicondyle	•11	•12-14



Clinical presentation

fall onto an outstretched hand

lateral elbow pain and swelling

may be subtle if fracture is minimally displaced

exam lacks the obvious deformity often seen with supracondylar fractures

swelling and tenderness are usually limited to the lateral side

lateral ecchymosis implies a tear in the aponeurosis of the brachioradialis and signals an unstable fracture

AP, lateral, and oblique views of elbow

internal oblique view most accurately shows fracture displacement because fracture is posterolateral

Treatment

Nonoperative

long arm casting 4-6wks

only if < 2 mm displacement in all views

medial cartilaginous hinge must remain intact

cast with elbow at approx 90 degrees as long as swelling is mild

weekly follow up and radiographs every week first 3 weeks, including internal oblique view

occasionally > 6 weeks of casting is needed

Operative

Closed reduction percutaneous pinning

1. **CRPP + 3-6 wks in above elbow cast**

   Fractures with 2 - 4 mm of displacement have intact articular cartilage and can be treated with CRPP

2. **Open reduction and fixation + 3-6 wks in above elbow cast**

   more than 4mm of displacement

   Open reduction (rather than closed) necessary to align the joint surface

   Joint incongruity

   Fracture non-union

A: Milch Type I

B: Milch Type II

Olecranon Fracture

Anatomy

- The olecranon and the proc. coronoideus form the Incisura trochlearis , which articulates with the trochlea of the distal humerus
- The intrinsic anatomy of this articulation allows for flexion/extension movement of the elbow joint and provides for stability of the elbow







Mechanism of injury

• Direct: Fall on the point of elbow or direct trauma to olecranon.

>> Typically results in a comminuted olecranon fracture.

• Indirect: Fall onto the outstretched upper extremity accompanied by a strong, sudden contraction of triceps.

>> Typically results in a transverse or oblique fracture.

• A combination (Both)



Clinical Presentation

• Patients typically present with the upper extremity supported by the contralateral hand with the elbow in relative flexion.

– Abrasions over olecranon or hand.

– Palpable defect at fracture site.

– Inability to extend the elbow actively against gravity.

– Ulnar nerve injury is possible, esp. with comminuted fractures

Radiographic Evaluation

True lateral radiograph:

Demonstrates: • extent of the fracture • degree of comminution • degree of articular surface involvement • displacement of the radial head, if present

AP view: This should be evaluated to exclude associated fractures or dislocations. –The distal humerus may obscure osseous details of the olecranon fracture.



Treatment

Nonoperative

- Immobilization in a long arm cast with the elbow in 45-90° of flexion is favored by many authors.
- In reliable patients a posterior splint or orthosis with gradual initiation of ROM after 5-7 days may be used.
- Follow-up radiographs should be obtained within 5-7 days after treatment to rule out fracture displacement.

Operative

.Rigid fixation is required

K-wire with tension band used primarily for .# transverse and short oblique

.Screw with tension band used also for the that

Plate fixation: • the best choice when comminution is present, to maintain ulnar length and alignment. when possible, interfragmentary .screws

Primary bone grafting should be considered to reduce the risk of nonunion

Treatment



Fragment Excision with Triceps Advancement

– Excision in patients with > 60% articular involvement yields poorer results.

– Patients must have an intact MCL, interosseous membrane, and DRUJ before excision, or instability will likely develop.

Radial Head Fracture

- Result from fall on outstretched hand
- May occur in elbow dislocation
- Swelling lat aspect
- Limited ROM
- Maximal tenderness over radial head

Immediate ortho referral criteria:

- fracture dislocation
- brachial artery or nerve injury
- 2mm displacement
- 1/3 of articulating surface
- Angulated > 30º
- Depressed > 3mm
- Severely comminuted

Radiograph

- AP and lat
- Fat pad may be only clue (occurs as a result of distension of the capsule by an intra-articular hemarthrosis)







ADIL RADIOLOGY

Mason-Johnston Classification of Radial Head Fractures

Classification	Description	Notes
Type 1	<2mm displacement	Conservative treatment
Type 2	>2 mm displacement	Conservative treatment if ROM preserved
Type 3	Comminuted	Surgical treatment
Type 4	Associated proximal radial dislocation	Surgical treatment

Treatment

long → above elbow
short → below

Non- displaced fracture:

- Immob in long-arm posterior splint with elbow flexed 90º.
- Ice and elevation for 48 hours
- Analgesia
- Forearm rotation out of splint 3-5 days
- 1 week – sling for comfort only
- Active ROM

Treatment

ORIF:

- Mason type II with mechanical block(displaced)
- Large fragment >2 mm
- Mason type III where ORIF feasible
- Mechanical block to motion
- Presence of other complex elbow injuries

Radial Head Replacement

- To prevent proximal migration of the radius
- Silicon implant poor outcome

>> SYNOVITIS

- Titanium/vitallium metallic implant of choice

Radial head excision

Indication:

- low demand, sedentary patients
- in a delayed setting for continued pain of an isolated radial head fracture

contraindication:

- in children
- presence of destabilizing injuries (essex-lopresti lesion, fracture dislocation elbow(mason type 4), monteggia)



- terrible triad of elbow(coronoid fracture,mcl deficiency)

Forearm Fractures

 Epidemiology

– Highest ratio of open to closed than any other fracture except the tibia

– More common in males than females, most likely secondary mva, contact sports, altercations, and falls

 Mechanism of Injury – Commonly associated with mva, direct trauma missile projectiles, and falls

Clinical Evaluation

– Patients typically present with gross deformity of the forearm and with pain, swelling, and loss of function at the hand

– Careful exam is essential, with specific assessment of radial, ulnar, and median nerves and radial and ulnar pulses

– Tense compartments, unremitting pain, and pain with passive motion should raise suspicion for compartment syndrome

Radiographic Evaluation

– AP and lateral radiographs of the forearm

– Don’t forget to examine and x-ray the elbow and wrist

Treatment

-- Conservative:

closed reduction + plaster fixation

In children, closed treatment is usually successful because the tough periosteum tends to guide and then control.

- Full length cast, from above elbow to metacarpal shaft

All adults unless the fragments are in close apposition >> Open reduction and internal fixation.

Complications:

- **Early:**
  - Compartment syndrome: from the fracture and operation.
  - Nerve injury: Posterior interosseous.
  - Vascular injury: radial or ulnar artery.
- **Late:**
  - Delayed union and non-union.
  - Malunion

Galeazzi's Fracture

- Radial shaft fracture at junction of middle and distal thirds with disruption of distal radio-ulnar joint
- Fall on extended pronated wrist
- Suspect if tenderness at distal radius and distal radial ulnar joint (DRUJ) disruption



Radiography

- Transverse or oblique fracture at junction of middle and distal thirds seen on AP view
- Widening of DRUJ on AP view
- Fracture of base of the ulnar styloid
- Radial shortening > 5mm
- Dislocation of radius relative to ulna on lat view

Treatment: Open Reduction Internal Fixation of the radial fracture plating

plate and screw

Monteggia's Fracture

- Fracture of ulnar shaft with dislocation of radial head
- Fall on outstretched, extended, and pronated elbow is usual mechanism
- Radial head may be palpated in antecubital fossa
- Radial nerve neuropraxia

Bado classification

- Type I- Anterior Dislocation of the radial head the most common (59%),
- Type II- Posterior/posterolateral dislocation of the radial head 26%
- Type III- Lateral/anterolateral dislocation of the radial head 5%
- Type IV- anterior dislocation of the radial head with fracture of radius and ulna at the same level 1%

Monteggia's Equivalents

- 1. Isolated radial head dislocation with plastic deformation of ulnar shaft
- 2. Fracture proximal ulna with radial neck fracture
- 3. Both bone proximal third fracture with radial fracture more proximal than ulna fracture



Radiology

Anterior dislocation of the radial head
Fracture of the proximal 1/3 of the ulna

Monteggia Fracture
EBM CONSULT
Radiocapitellar Line
Humerus
Capitellum
Medial Epicondyle
Radial Head
Ulna
Radius

Treatment

The elbow is immobilized in a long-arm cast for 4 weeks in 110 to 120 degrees of flexion with the forearm in neutral rotation.

A short-arm cast is used for an additional 4 weeks while early range of motion at the elbow and forearm is begun.

OPERATIVE TREATMENT

Two indications for operative treatment of type I fracture dislocations

1). failure of ulnar reduction

2). failure of radial head reduction.

3) All adult fractures.

Distal Radius Fractures

 Epidemiology

– Most common fractures of the upper extremity

– Common in younger and older patients.

Usually >> fall on out stretched hand

 Mechanism of Injury

– Most commonly a fall on an outstretched extremity with the wrist in dorsiflexion – High energy injuries may result in significantly displaced, highly unstable fractures

Colles' fracture

Fr. distal 1 inch of the radius, with or without avulsion of the ulnar styloid.

Common in old age due common in old age due to osteoporosis.

Most common of all fractures in older people

Dinner-fork deformity

Distal Radius Fractures



– **Colles** Fracture most common

- Combination of intra and extra articular fractures of the distal radius with dorsal angulation (apex volar), dorsal displacement, radial shift, and radial shortenting
- Most common distal radius fracture caused by fall on outstretched hand

– **Smith** Fracture (Reverse Colles)

- Fracture with volar angulation (apex dorsal) from a fall on a flexed wrist



– **Barton** Fracture

- Fracture with dorsal or volar rim displaced with the hand and carpus

– **Radial Styloid Fracture (Chauffeur Fracture)** Hutchinson fractures and backfire fractures.

- Avulsion fracture with extrinsic ligaments attached to the fragment
- Mechanism of injury is compression of the scaphoid against the styloid



LearningRadiology.com (C)
All Rights Reserved
LearningRadiology.com (C)
All Rights Reserved
RS

Treatment

– Displaced fractures require and attempt at reduction

- Hematoma block-10ccs of lidocaine or a mix of lidocaine and marcaine in the fracture site
- Hang the wrist in finger traps with a traction weight
- Reproduce the fracture mechanism and reduce the fracture
- Place in sugar tong splint

– Operative Management

- For the treatment of intra-articular, unstable, mal-reduced fractures.
- As always, open fractures must go to the OR.

Treatment

Barton is unstable after reduction unlike Smith

-Displaced fractures require and attempt at reduction.

- Hematoma block-10ccs of lidocaine or a mix of lidocaine and marcaine

-Hang the wrist in fingertraps with a traction weight

- Reproduce the fracture mechanism and reduce the fracture

-Place in sugar tong splint

Burton

Bolton

Dinner fork
distal head fracture

nightstick fracture


Subcondylar


Normal

Radial head luxation
radiocapitellar line
radius
capitellum

capitellum
radiocapitellar line


Subcondylar


Radial dislocation

Supracondylar

A
B

How would you manage this patient?


R
BAM

Thank you for your attention

Hip joint and peri-trochanteric trauma

Presented By

Dr. Karrar Kareem

Orthopedic Surgeon

Anatomy

- The hip joint has a ball-and- socket configuration; synovial articulation between the head of the femur and the acetabulum of the pelvis bone.
- 40% of the femoral head is covered by the bony acetabulum at any position of hip motion. The effect of the labrum is to deepen the acetabulum and increase the stability of the joint by 10% .


Anatomy of the Hip and Acetabular Labrum
Chondral surfaces of the femoral head and acetabulum
Femoral head
Acetabulum
Joint capsular ligament
Acetabular labrum
Greater trochanter
Lateral (side) view with femur removed of right hip and acetabulum
Anterior (front) cut-away view of right hip and acetabulum

The joint is supplemented by much stronger ligamentous condensations iliofemoral, pubofemoral, and ischiofemoral ligaments that run in a spiral fashion, preventing excessive hip extension.



Main vascular supply is from the lateral and medial femoral circumflex arteries, branches of the profound femoral artery.

An extracapsular vascular ring is formed at the base of the femoral neck with ascending cervical branches that pierce the hip joint at the level



Classification

Depending upon the position of the head with respect to the acetabulum, hip dislocations are classified as:

- Posterior dislocations: Commonest and is seen in 80-90 percent of the cases.
- Anterior dislocations: Seen in 10-15 percent.
- Central dislocations: Relatively rare

Posterior Dislocation
Also known as “dashboard injury”
They result from trauma to the flexed knee, with the hip in varying degrees of flexion.



THOMPSON AND EPSTEIN CLASSIFICATION

- Type I: With or without minor fracture.
- Type II: With a large single fracture of the posterior acetabular rim.
- Type III: With communition of the rim of the acetabulum with or without a major fragment.
- Type IV: With fracture of the acetabular floor.
- Type V: With fracture of the femoral head.



ANTERIOR DISLOCATION

•Hyperextension force against an abducted leg that levers head out of acetabulum.

Femoral head dislocated anterior to acetabulum In RTA's, when the knee strikes the dashboard with the thigh abducted.



The femoral head is medial inferior go the actabulum

CENTRAL DISLOCATION

This is the least common and most difficult of all dislocations of the hip joint.

Mechanism of Injury- It could be due to direct blow on the greater trochanter as in the case of RTA or fall on the sides

It is invariably associated with the fractures of the acetabulum and this is what makes it a very difficult problem to treat



Management History and Evaluation

Physical Examination

Lower limb is flexed, adducted and internally rotated.

Shortening and reverse in anterior dislocation

Femoral head palpated post.

- Narthes sign (i.e. Difficulty to palpate femoral pulse due to backward migration of femoral head).

Movement Painful limitation of all hip movements.

Usually posterior

Signs of sciatic nerve injury:

Loss of sensation in posterior leg and foot

Loss of dorsiflexion (peroneal branch) or plantar flexion (tibial branch)

Loss of deep tendon reflexes at the ankle S1,2

Signs of femoral nerve injury:

Usually anterior

Loss of sensation over the thigh

Weakness of the quadriceps

Loss of deep tendon reflexes at knee L3, 4

TREATMENT

All hip dislocations are emergencies and need to be reduced

To prevent troublesome late complications like AVN and traumatic degenerative hip.

Methods of Closed Reduction

Allis method

Bigelow method

Classical Watson Jones method

Stimson's gravity method

Whistler's technique(over-under)

Captain Morgan technique (Nobel method)

Allis Method

The patient is placed supine the surgeon standing above the patient on the stretcher or table •. Initially, the surgeon applies in- line traction while the assistant applies counter traction by stabilizing the patient's pelvis.
•While increasing the traction force, the surgeon should slowly increase the degree of flexion to approximately 70 degrees. • Gentle rotational motions of hip as well as slight adduction will often help the femoral head to clear the lip of the acetabulum. • A lateral force to the proximal thigh may assist in reduction. An audible "clunk" is a sign of a successful closed reduction.



STIMSONS GRAVITY METHOD

this is the reverse allis method of reduction. The steps are as follows: • patient is prone • patient is brought to the edge of the table. • An assistant stabilizes the pelvis by applying downward pressure over the sacrum • the affected hip and knees are flexed to 90 degrees. • Downward pressure is applied on the flexed knee. • To facilitate the reduction, gentle rotations needs to be done.



Non-operative Treatment

If hip stable after reduction:

- Maintain patient comfort
- skin traction , analgesia
- Avoid Adduction, Internal Rotation.
- No flexion > 60
- Early mobilization usually few days to 2 weeks.
- Repeat x-rays before allowing full weight-bearing.

Indications for Open Reduction:

- Failed closed reduction.
- Failed stability test.
- Big posterior lip fragment.
- Bone fragment within the acetabulum.
- Fracture of the femoral head.
- Sciatic nerve palsy.

Postoperative Treatment

- Skeletal traction (10-15 lbs) with the hip in slight abduction and extension.
- Within 3-5 days, gentle active and passive exercises in traction are begun.
- Traction to be maintained for 6-8 weeks.
- Later protected weight bearing is allowed

Up to 3 months

Complications

- Myositis ossificans (2%)
- Traumatic osteoarthritis due to avascular necrosis (35%)
- Sciatic nerve injury: 10-13% , it is 3 times more common in fracture dislocation than simple dislocation

Commonly when the hip is less flexed

- Irreducible dislocation (31%)

Femoral Head Fractures

It usually comes as an association



Extracapsular arterial ring:

- medial circumflex femoral artery (main supply to the head) from profunda femoris
- lateral circumflex femoral artery
- ascending cervical branches
- artery to the ligamentum teres

from the obturator artery or MCFA

supplies peri-foveal area

-- Anterior hip dislocation.

-- Posterior hip dislocation.

location and size of the fracture fragment and degree of comminution depend on the position of the hip at the time of dislocation.

Pipkin Classification

- Type I Fx below fovea/ligamentum (small) Does not involve the weightbearing portion of the femoral head
- Type II Fx above fovea/ ligamentum (larger) Involves the weightbearing portion of the femoral head
- Type III Type I or II with associated femoral neck fx High incidence of AVN
- Type IV Type I or II with associated acetabular fx (usually posterior wall fracture)



Presentation and Radiographs



- **History**
  - frontal impact MVA with knee striking dashboard
  - fall from height
- **Symptoms**
  - localized hip pain
  - unable to bear weight
  - other symptoms associated with impact
- **Physical exam**

shortened lower limb and others

- AP pelvis, lateral hip and Judet views
- both pre-reduction and post-reduction inlet and outlet views
- CT scan for post reduction to evaluate concentric reduction loose bodies in the joint acetabular fracture femoral head or neck fracture

Non-operative and operative

- Hip reduction

>> acute dislocations and reduce hip dislocation within 6 hours

>> obtain post reduction CT

- TDWB x 4-6 weeks, restrict adduction and internal rotation in:

  - Pipkin I
  - undisplaced Pipkin II with < 1mm step off
  - no interposed fragments
  - stable hip joint

- Operative --ORIF indications

  - Pipkin II with > 1mm step off
  - if performing removal of loose bodies in the joint
  - associated neck or acetabular fx (Pipkin type III and IV)
  - polytrauma
  - irreducible fracture-dislocation
  - Pipkin IV

Arthroplasty

- indicationsPipkin I, II (displaced), III, and IV in older patients
- Fractures that are significantly displaced, osteoporotic or comminuted

Total if associated with other
Partial if low demand and isolated fractures



Femoral neck fracture

Mechanism:

- high energy in young patients
- low energy falls in older patients

Osteology:



- normal neck shaft-angle 130 +/- 7 degrees
- normal anteversion 10 +/- 7 degrees

Factor delay femoral fracture healing in femoral neck

- intra articular because synovial fluid prevent formation of hematoma which is associated with callus formation

-thin periosteum

-poor blood supply

Blood supply to femoral head and neck:

- major contributor is medial femoral circumflex (lateral epiphyseal artery)
- some contribution to anterior and inferior head from lateral femoral circumflex
- some contribution from inferior gluteal artery
- small and insignificant supply from artery of ligamentum teres

Radiographs

- Recommended views obtain AP pelvis and cross-table lateral, and full length femur film of ipsilateral side
- Garden classification is based on AP pelvis
- CT helpful in determining displacement and degree of comminution in some patients



Broken line indicate operative in children unlike adult

Classification (PAUWEL , GARDEN)



Fig-18.9 Pauwel's classification

Easily displaced

Valgus impaction



Non-operative and operative

All are surgical indications

observation alone

- indications : may be considered in some patients who are non-ambulators, have minimal pain, and who are at high risk for surgical intervention

Surgical Rx:

cannulated screw fixation

Sliding Hip Screw Or Cephalomedullary Nail

Just in basal cervical fracture

Hemiarthroplasty and arthroplasty

Cannulated screw fixation

Adult
Undisplaced// colsed fixation (cannulated screw in v shape configuration
Dispaced / open reductions + cannulated screw fixation

Elderly
Active - displaced = total hip replacement
- undisplaced = cannulated screw fixation

Inactive /p- displaced = partial hip replacement



➢ Indications

- nondisplaced transcervical fx
- Garden I and II fracture patterns in the physiologically elderly
- displaced transcervical fx in young patient

considered a surgical emergency achieve reduction to limit vascular insult

>> reduction must be anatomic, so open if necessary



Sliding Hip Screw Or Cephalomedullary Nail

Also dynamic hip screw



- **Indications**
- basicervical fracture
- vertical fracture pattern in a young patient
- biomechanically superior to cannulated screws
- consider placement of additional cannulated screw above sliding hip screw to prevent rotation



Hemiarthroplasty and arthroplasty

- **Hemiarthroplasty:**

Indications debilitated elderly patients metabolic bone disease

- **Total hip arthoplasty:**

Indications older active patients patients with preexisting hip osteoarthritis

 more predictable pain relief and better functional outcome than hemiarthroplasty

 arthroplasty for Garden III and IV in patient < 85 years

INTERTROCHANTERIC FRACTURES

Extracapsular fractures of the proximal femur between the greater and lesser trochanters





INTERTROCHANTERIC FRACTURES

Mechanisms:

elderly low energy falls in osteoporotic patients young high energy trauma

Deforming muscle forces will usually produce [1]shortening, [2]external rotation and [3]varus position at the fracture

>> Occur in the region between the greater and lesser trochanters of the proximal femur, occasionally extending into the subtrochanteric region

>> Since they occur in cancellous bone with abundant blood supply – no problems of non-union and osteonecrosis



Deforming Forces



INTERTROCHANTERIC FRACTURES

90% of intertrochanteric fractures in the elderly result from a simple fall

The tendency to fall increases with patient age and is exacerbated by several factors, including

(poor vision, decreased muscle power, labile blood pressure, decreased reflexes, vascular disease, and coexisting musculoskeletal pathology.)

Most fractures result from a direct impact to the greater trochanter area

 Low energy falls from a standing height – approximately 90% of community hip fractures in patients more than 50 years of age with a higher proportion of women

Pre-surgery workup

Lab investigations

Viral screening

Imaging

Echo study and ECG

Classification Kyle's Classification



TYPE 1
Undisplaced
Uncomminuted



TYPE 2
Displaced
Minimal comminuted
Lesser trochanter fracture
Varus



TYPE 3
Displaced
Greater trochanter fracture
Comminuted
Varus



TYPE 4
Severely comminuted
Subtrochanter extension
(Also reverse oblique)



Sign of instability
-reverse oblique fracture line
-inferiomedial aspect comminuted (varus)
-cyst formation in fracture site

Treatment

These fractures are almost always treated by early internal fixation

Non-Operative Rx:

Buck's traction or extension

Russell skeletal traction

Balanced traction inThomas splint

Plaster spica immobilization

Derotation boot

not because they fail to unite with conservative treatment (they unite quite readily), but (1) to obtain the best possible position and (2) to get the patient up and walking as soon as possible.

Complication of Non-Operative Mx:

Decubiti, UTI, joint contractures, pneumonia, and thromboembolic complications, resulting in a high mortality rate.

In addition, fracture healing is generally accompanied by varus deformity and shortening because of the inability of traction to effectively counteract the deforming muscular forces

Neck=nonunion
Inter trochanter =malunion

Operative Rx

As soon as the general condition of this patient is under control, internal fixation should be carried out.

 The goal of surgical treatment is strong, stable fixation of the fractured fragments

Factors affecting ORIF:

Bone quality Fracture geometry Reduction Implant design Implant placement

Choices of treatment in intra trochanter

-intra medullary nail(gamma nail) =top choice

-locked plate- 2nd choice

-open with dynamic hip screw =3rd choice

-blade plate= last choice

ORIF CHOICES



Subtrochanteric fractures

just below the level of the lesser trochanter.

10 -30% Of Hip Fractures

- Bimodal Age Distribution
  - **20-40yrs**
- High Energy
- RTA, Fall From Height, Gsw (10%)
  - **>60yrs**
- Low Energy
- Simple Fall, Osteoporotic, Pathological (17-35%)
- Cann. Screws Below Lesser Troch
- Bisphosphonate Rx

Between lesser trochanter and 5cm distal

- Stresses
  - Posteromedial – Lateral

Compression on the medial, tension on the lateral

  - compressive – tensile
  - 6 x BW
- Cortico-cancellous junction
  - Vascularity
  - Healing

Deforming forces

Proximal fragment

– Flexed – iliopsoas

– Ext rot – iliopsoas + short ext rots

– Abducted – gluteus med + min

• Distal fragment

– Adducted/ varus – adductors – Shortened – quads/ hamstrings

3 deformety forces of proximal segment
- flexion(iliopsoas)
-abduction(gluteus medius+ minimus)
-external rotation (piriformis)



History

- Pain, NWB, deformity of thigh
- Open/ closed injury
- Neurovascular status
- Full ATLS – Haemorrhagic shock

– Cranial, thoracic, abdominal injuries (Waddell's triad)

Russell-Taylor Classification



Treatment (conservative vs operative)

All patient should undergo surgery unless unfit for surgery

Open reduction and internal fixation is the treatment of choice.

Unfit for surgery, open #s

• Skeletal traction

Hamilton-Russell, Perkins, 90-90 position

<5 varus/valgus, >25% cortical apposition, <1cm shortening

– No axial rotation• Weekly x-rays + adjustments

• Traction until

signs of union

(8-12 weeks)











Surgery





Key issues

– Medial cortical buttress

– Reduction (esp varus deformity)

• Extramedullary

– DHS, Medoff Sliding Plate, DCS, blade plate, locking plates

• Intramedullary – Centromedullary – Cephalomedullary – Short vs long

First line

Intramedullary
Blade plate
Locked plat
Dynamic hip screw

Long

Short for inter trochanter

Complications

- Infection
- VTE
- Implant failure,
  - Varus malreduction,
  - Screw placement in femoral head,
- Malunion
  - Shortening – Rotational deformity – Varus
- Nonunion (0-8%)
  - Significant pain >6 months with the inability to FWB – Stable fixation - autogenous bone grafting
  - Exchange nailing with over-reaming



Femur Shaft Fractures

Anatomy

Anteriolateral

Long tubular bone, anterior bowed forward and has oblique course from the neck to distal end.

Femoral shaft fracture is a fracture of the femoral diaphysis occurring between 5 cm distal to the lesser trochanter and 5 cm proximal to the adductor tubercle







Compartments of thigh Muscle of the thigh are arranged in three Compartments

1-anterior compartment of thigh contains the sartorius and the four large quadriceps

2-medial compartment of thigh (gracilis ,pectineus, adductor longus, adductor brevis, adductor magnus, and obturator externus)

3-posterior compartment of thigh contain three large muscle termed the ‘”hamstring” which provide the blood supply to femur



Epidemiology

-Common injury due to major violent trauma femur fracture more common in people < 25 yr or >65 yr

-Motor vehicle, motorcycle, auto-pedestrian, fall from height, and gunshot wound

Mechanism Of Injury

• In Young Adults, almost always the result of high-energy trauma

– Motor vehicle accident

– Gunshot injury, or

– Fall from a height

• Pathologic fractures, especially in the elderly, commonly occur following a trivial fall

• Stress fractures occur mainly in military recruits or runners

Signs and symptoms

Diffuse pain or ache, and tenderness and swelling in the thigh or groin.

Bleeding and bruising in the thigh (uncommon).

Non ambulatory with pain –Variable gross deformity of thigh –Swelling, –Shortening of the affected extremity

Weakness

pallor

Cont…

A careful assessment of hemodynamic stability is essential

Average expected Blood loss of 750-1500ml

- Thorough examination of the ipsilateral hip and knee should be performed
- Knee ligament injuries are common, however, and need to be assessed after fracture fixation

Associated Injuries

- Ipsilateral femur neck, intertroch, distal femur #s
- Patella, tibia, acetabular, pelvic ring #s
- Soft tissue injuries of knee
- Thoracic & abdominal injuries(5-15%)

X-ray

X-ray Will confirm the diagnosis and establish the sites ,line ,extent and displacement

- AP and Lateral views of the femur, hip, and knee
- AP view of the pelvis should be obtained
- Look for evidence of an associated femoral neck or intertrochanteric fracture, knee injuries





Management

A) > 6 m , support.

B) 6m - 5 yrs , cast

C) 5 - 13 yrs :

BW <55 kg , transverse fracture , short oblique flexible nail .
BW >55 kg , spiral fracture , long oblique plate and screw .

D) 13 - 16 yrs , trochantric nail ,

Intramedullary nailing is the method of choice for most femoral shaft fractures.

Flexible intra nail in:
-Less than 55
-transverse or low oblique

Plating

Indications for plating

- Patients wit extremely narrow medullary canal
- Fractures around or adjacent previous malunion
- Fractures extending proximally or distally into the pertrochanteric or metaphyseal region
- Ipsilateral neck fractures



External Fixation

- Severe soft tissue injuries
- Contamination
- Associated vascular injuries
- Polytrauma-temporary method



Complications

1-GENERAL

Blood loss, shock ,fat embolism, and acut respiratory distress are common in high- energy injuries

2-Vascular injury

3-Nerve injury-iatrogenic(femoral, sciatic, pudendal, peroneal)

4-Thromboembolism

5-Compartment syndrome(1-2%)

6-Infection

7-Delayed union and non-union

8-Joint stiffness, knee & hip pain

9-Heterotrophic ossification

THANK YOU

Lower limb trauma...

Presented by

Dr. Karrar Kareem

Orthopedic surgeon

Distal femur fracture

Definition Fractures of the thighbone that occur just above the knee joint are called distal femur fractures.

Anatomy

Basic anatomy of femur - only bone in the thigh.

- It is classed as a long bone, and is the longest bone in the body.
- The main function of the femur is to transmit forces from the tibia to the hip joint.

Articulate proximally with hip bone forming hip joint and distally with tibia & patella forming knee joint

Divided into three areas: proximal, shaft and distal

tricompartmental

The distal end is characterised by the presence of the medial and lateral condyles, which articulate with the tibia and patella, forming the knee joint.

Medial and lateral condyles – Rounded areas at the end of the femur. The posterior and inferior surfaces articulate with the tibia and menisci of the knee, while the anterior surface articulates with the patella.

Medial and lateral epicondyles – Bony elevations on the non-articular areas of the condyles. They are the area of attachment of some muscles and the collateral ligaments of the knee joint.


Medial epidcondyle
Lateral epicondyle
Medial condyle
Lateral condyle
Intercondylar fossa


Adductor tubercle
Lateral epicondyle
Medial epicondyle
Lateral condyle
Medial condyle

Path mechanics

When the distal femur breaks, both the hamstrings and quadriceps muscles tend to contract and shorten.

When this happens the bone fragments change position and become difficult to line up with a cast.

gastrocnemius: extends distal fragment (apex posterior)

adductor Magnus: leads to distal femoral Varus




Quadriceps
Hamstrings


Plantaris
Heads of the gastrocnemius
Soleus
Calcaneal tendon
© teachmeanatomy

Investigations

X-ray .. obtain standard AP and Lateral

CT .. obtain with frontal and sagittal reconstructions useful for establishing intra-articular involvement

Angiography : indicated when diminished distal pulses after gross alignment restored, consider if associated with knee dislocation

Hoffa fracture

Is a type of supracondylar distal femoral fracture and is characterized by an associated fracture component in the coronal plane.

- Hoffa fractures are intra-articular and are characterised by a fracture in the coronal plane. Always displaced
- Hoffa fragments are more commonly unicondylar and usually originate from the lateral femoral condyle.

They can be occasionally bicondylar.

PODCAST PEARLS

Treatment

Previously but now it is rare

Non - Operative

Skeletal traction

Casting and bracing for 6 weeks : indications (rare) >> - non displaced fractures - non ambulatory patient - patient with significant comorbidities

Operative

External fixation temporizing measure until soft tissues permit internal fixation, or until patient is stable
avoid pin placement in area of planned plate placement if possible

ORIF indications : 1- displaced fracture 2- intra-articular fracture 3- nonunion - goals : 1- need anatomic reduction of joint 2- stable fixation of articular component to shaft to permit early motion 3- preserve vascularity

Retrograde IM Nail

All fractures are fixed by plate and screw except isolated suprachondylar fracture in which IM nail is used


Pins
Femur
Pins
Tibia

Postoperative

Early ROM of knee important

non-weight bearing or toe touch weight-bearing for 6-8 weeks, up to 10-12 weeks if comminuted * quadriceps and hamstring strength exercises



Anatomy

- The **knee joint:**

Is a hinge **type** synovial **joint**, which mainly allows for flexion and extension (and a small degree of medial and lateral rotation)

- **Ligaments**

PCL, ACL, LCL, MCL, and PLC are all at risk for injury

main stabilizers of the knee given the limited stability afforded by the bony articulations

- **Blood supply**

popliteal artery injuries occur often due to tethering at the popliteal fossa

  - proximal - fibrous tunnel at the adductor hiatus
  - distal - fibrous tunnel at soleus muscle

geniculate arteries may provide collateral flow and palpable pulses masking a limb-threatening vascular injury

Blood Supply of the Knee Joint

- **Femoral**

  Descending genicular – articular and saphenous

- **Popliteal**

  Superior medial genicular, middle genicular, inferior medial genicular, superior lateral and inferior lateral genicular

- **Tibial**

  Anterior and posterior tibial recurrents

- Also, anastamosis from descending branch of lateral circumflex femoral aa
- **Middle genicular branch** sends fibers inside the joint to supply cruciate ligament, synovial membrane and periphery of menisci



NERVE SUPPLY

Apply Hilton's law

The nerve supplying a joint also supplies both the:

- Muscles that move the joint &
- Skin covering the articular insertion (area of action) of those muscles

- Femoral
- Obturator
- Sciatic
- Nerve to quadratus femoris

These nerves also supply knee joint & therefore pain arising in the hip joint may be referred to the knee

Knee dislocations

Knee dislocations are traumatic injuries characterized by a high rate of vascular injury

- treatment is generally emergent reduction and assessment of limb perfusion

Epidemiology

- incidence
  - 0.02% of orthopedic injuries
  - likely underreported as approximately 50% self-reduce and are misdiagnosed
- demographics
  - 4:1 male to female
- location
  - tibiofemoral articulation (knee joint)
- risk factors
  - morbid obesity is a risk factor for "ultra-low energy" knee dislocations with activities of daily living

Mechanism of injury

- high-energy vs low energy
- hyperextension injury leads to anterior dislocations
- posteriorly directed force across the proximal tibia (dashboard injuries) leads to posterior dislocations

Associated injuries:

Because it is lateral and tethered

- Vascular injury
- Nerve injury >> peroneal 25%, tibial n. less common
- Fractures in about 60%
- Soft tissue injury



Classification

- **Kennedy classification**

Direction of displacement of the tibia

- anterior 30-50%
- posterior 30-40%
- lateral 13%
- medial 3%
- rotational 4%

Schenck Classification

- **KD I** injury to the ACL or PC
- **KD II** Injury to ACL and PCL
- **KD III** injury to ACL, PCL, and PMC or PLC
- **KD IV** Injury to ACL, PCL, PMC, and PLC
- **KD V** Multiligamentous injury with periarticular fracture

Physical exam

Appearance

No obviuos >> 50% spontaneously reduce before arrival to ED

Obviuos >>reduce immediately, especially if absent pulses

Vascular exam

if pulses are present and normal

If ABI >0.9 >> then monitor with serial examination

If ABI <0.9 >> perform an arterial duplex ultrasound or CT angiography

if arterial injury confirmed then consult vascular surgery

Neurologic exam:

Sensory and motor function of peroneal and tibial nerve as nerve deficits often occur concomitantly with vascular injuries



Stability:

Assess ACL, PCL, MCL, LCL, and PLC

diagnosis based on instability

Imaging

Pre and post-reduction AP and lateral of the knee

- look for asymmetric or irregular joint space
- look for avulsion fxs (Segond sign - lateral tibial condyle avulsion fx)
- Osteochondral defects

CT

MRI



Treatment

Nonoperative

>> Emergent closed reduction followed by vascular assessment/consult

>> Immobilization as definitive management

Operative

Emergent open reduction

> irreducible knee

> posterolateral dislocation

> open fracture-dislocation

> obesity

> compartment syndrome

> vascular injury

external fixation

delayed ligamentous reconstruction/repair

Complications of knee dislocation

Vascular compromise

>> 40-50% in anterior or posterior dislocations

>> KD IV injuries have the highest

Rx >> emergent vascular repair and prophylactic fasciotomies

Stiffness (arthrofibrosis)

most common complication (38%)

more common in delayed mobilization

Mx by early reconstruction and motion, arthroscopic lysis of adhesion, and manipulation under anesthesia

Laxity and instability:

37% of some instability, however, redislocation is uncommon

Peroneal nerve injury

25% occurrence of a peroneal nerve injury

50% recover partially

posterolateral dislocations

Rx >> AFO to prevent equinus, neurolysis or exploration later on, tendon transfers if chronic nerve palsy persists

Ankle foot orthoses

Patella - Anatomy

• Largest sesamoid bone

• Plays an important role in the biomechanics of the knee.

• Very hard & triangular-shaped bone

• Situated in an exposed position in front of the knee joint

• separated from the skin by subcutaneous bursa.

• Patella Surfaces ▫ Ant. ▫ Post ▫ Lat. & med.

Patella borders ▫ Base ▫ Med and Lat. ▫ Apex • Articulation

Post. surface central portion, is covered with a layer of hyaline cartilage.

• Articular cartilage of the patella is the thickest in the body (up to 7-mm thick)

Improves the efficiency of extension during the last 30° of knee extension.

Patella fracture

- Direct impact
- Indirect trauma in which a severe pull of the patellar tendon occurs when the knee if semi-flexed (Commonest)

S/S
- hemorrhage which results in significant swelling
- pain
- POT over Patella
- extreme pain with weight bearing/movement

Functions

- Guide for the quadriceps or patellar tendon
- ↓ Friction of the quadriceps mechanism
- Acts as a shield for the femoral condyles
- Improves appearance of the knee.

Classification

Undisplaced

- Transverse fracture (80%)
- Vertical fracture
- Comminuted fracture

Displaced

- Transverse (85 %)
- Vertical fracture
- Comminuted fracture
- Osteochondral fracture



Investigations

X – ray

- AP view
- lateral view
- Skyline view

CT scan

Bone scan

MR



Fig-20.5 Skyline view of patella

Treatment

Non operative

–For non displaced fracture cylinder cast: extending from the groin to just above the malleoli for 4 to 6 weeks.

–Followed by physiotherapy-quadriceps strengthening exercise

Operative

• Tension band wiring.

• Patellectomy Rarely indicated

1. Partial:for proximal pole fracture; major fragment is preserved

2. Complete: for comminuted fractures. >> Knee should be immobilized for 3 to 6 weeks in a long leg cast at 10degrees flexion for both partial and complete patellectomy.

• Open reduction and internal fixation for transverse fracture

Complications

• Refracture • Non union • AVN • OA • Stiffness • Patellar instability Incomplete extension



Lateral Bipartite Patella - AP and Skyview

Tibial plateau fractures

The tibial plateau is the proximal end of the tibia including the metaphyseal and epiphyseal regions as well as the articular surfaces made up of hyaline cartilage.

- AO defines tibial plateau as the metaphysis to a distal distance equal to the width of the proximal tibia at the joint line.

Distinction between medial and lateral condyles

Medial: Slightly concave shape, Larger in both width and length,Cartilage thickness ~ 3 mm

Lateral: Convex, 2-3 mm superior (proximal) to the medial, Cartilage thickness ~ 4 mm


Apex,
Head,
Neck
Soleal line




Interior superior view of tibia
Oblique popliteal lig.
Posterior meniscofemoral lig.
Posterior cruciate lig.
collateral lig.
Bursa
Tibial collateral lig.
Medial meniscus
Synovial membrane
Joint capsule
Anterior cruciate lig.
Patellar lig.
Anterior aspect

n extension: posterior view
Posterior cruciate lig.
Anterior cruciate lig.
Posterior meniscofemoral lig.
Medial condyle of femur
Lateral condyle of femur
Medial meniscus
Popliteus tendon
Lateral collateral lig.
Medial collateral lig.
Lateral meniscus
Head of fibula

In flexion: anterior view
Medial condyle of femur
Medial collateral lig.
Medial meniscus
Transverse lig.
Tibial tuberosity


aPPTA 81°(83)
aAD.

Mechanism of injury and Classification

Force directed medially (valgus deformity) or laterally (varus deformity) or both.

Axial compressive force.

Both axial force and force from the side.

• Shatzker classification – Six types

Type I: Split-wedge

Type II: Split +depression commonest

Type III: pure depression

Type IV: Split fracture of medial plateau

Type V: bicondylar fracture

Type VI: total disconnection from the diaphysis



Type 4,5,6 even if they not displaced they need surgery
Plate and screw

Hohl moore and Shatzker



ASSOCIATED INJURIES

- 90% of these fractures asssociated with soft tissue injuries
- Meniscal tears occurs in 50% of these fractures
- Associated ligamentous injuries (cruciate or collateral) occur in 30% of these fractures
- Others-: common Peroneal nerve
  Popliteal artery injury

Evaluation

- X ray AP view Lateral view
- CT with3D reconstruction - Better visualisation - Preoperative planning
- MRI Useful for evaluating injuries of menisci, cruciate & collateral ligaments and soft tissue envelope
- Arteriography for any vascular injury in question







Management

• Non-operative management: – Indicated for non-displaced or minimally displaced fractures

• Method:

– Protected weight bearing and early range-of-knee motion in a hinged fracture brace.

– Isometric quadriceps exercises and progressive passive, active-assisted, and active range-of-knee motion exercises.

– Partial-weight bearing (30-40 Ib) for 8 to 12 weeks with progression to full weight bearing.

Operative treatment

Accepted range of articular depression varies from < 2 mm to 1 cm – Instability > 10 degrees of nearly extended knee compared to the contralateral side – Open fractures – Associated compartment syndrome – Associated vascular injury

Goals of treatment: – reconstruction of the articular surface – re-establishment of tibial alignment • Treatment involves reducing and buttressing of elevated articular segments with bone graft • Soft tissue reconstruction including menisci and ligaments • Spanning external fixator as a temporizing measure in patients with high-energy injuries or significant soft tissue injury. • Arthroscopy

Criteria for conservative manegment

1-Gapping less than 1cm 5 MM 5
2-Articular Step less than 2mm

Type 4,5,6 even if they not displaced they need surgical manegment
Plate and screw

L

A

B

C

Tibial Shaft Fracture

Most common long bone fracture - 492,000 fractures/year

- Most common open fracture
- Significant cost - 569,000 hospital days

- Major cause of disability

- Significant complications

- 50,000 nonunions/year

Assessment

- History: velocity of injury
- Clinical examination
- General ATLS (advanced trauma life support)
- Local soft tissues
- Neurovascular

Assessment

- X-rays: A/P and lateral
- Special investigations:

Pressure monitoring

Angiography

CT (computed tomography, never immediate)



Nonoperative treatment

•Nonoperative treatment does NOT mean no treatment

• Closed reduction and plaster of Paris application achieve good results

• Nonoperative treatment is difficult and demanding

• Plaster can prevent lateral shift

• Plaster can prevent angulation

• Plaster can control rotation

• Plaster can **NOT** prevent shortening

Indications:

• Children

• Undisplaced fractures

• "Stable" reduced fractures

• Contraindication for surgery

Operative treatment

Tibial fixation options

- Plate
- Ex Fix
- IMN



Ankle fractures

Ankle is a three bone joint composed of the tibia, fibula an talus

- Talus articulates with the tibial plafond superiorly

Itsconsidered saddle-shaped with the dome itself is wider anteriorly than posteriorly



Imaging

An initial evaluation of the radiograph should 1st focus on

- Tibiotalar articulation and access for fibular shortening
- Widening of joint space
- Malrotation of fibula
- Talar tilt

Identifies fractures of

- malleoli
- distal tibia/fibula
- plafond
- talar dome
- body and lateral process of talus
- calcaneous

Danis Weber Classification





Normal



Weber A


Some fibres of the syndesmosis may be torn but the tibiofibular articulation is not widened
Often associated with medial malleolus fracture or deltoid ligament disruption

Weber B

Weber C

Thank you

Injuries of the ankle

Functional anatomy:

- The ankle act to support & propel the body ….. During running & jumping , loads well in excess of 10 times body weight are transmitted through the ankle & foot ,so if loading is excessive or repeated … lead to ankle & foot injury
- The ankle is close fitting hinge joint . made by distal end of tibia and fibula(mortise) and upward projection of talus(tenon) The mortise bones are held together as syndesmosis by distal tibiofibular lig. and iterossess membrane

The ankle moves only in one plane (flexion / extension).







More than 90% of injuries is in the lateral

1. The lateral ligament is weaker than the medial ligament.
2. The lateral ligament resists inversion.

the lateral collateral ligaments consist of:

-anterior talofibular ligament.

-posterior talofibular ligament.

-calcaneo fibular ligament.

usually the anterior talofibular, or both this and the calcaneofibular ligament; only in the most severe injuries is the posterior talofibular ligament torn.

medial collateral(deltoid) ligament:

- superficial
- deep





Ankle ligament injuries:

- A sudden twist of the ankle tenses the structures around the joint; this may amount to no more than painful wrenching of the soft tissue--------sprained ankle.

  If more severe force is applied :

- partial tear (most of the ligament remain intact --- once it heals it is able to support the weight of body.
- Complete tear: the ligament may still heal but never regain its original form ----------unstable.



Injuries of the lateral ligaments:

- 90 %of ankle injuries
- Mostly talofibular ligament.
- Mechanism of injury : inversion + planterflexion



Clinical features:

- -If the patient is able to walk ,& the bruising is only faint ,, it is probably sprain
- If the bruising is marked & patient unable to put weight on the foot ,,, suggest severe injury
- tenderness ; max just distal and ant. To lateral malleolus.
- - passive inversion --------painful.

- ***x-ray:*** AP,LAT,MORTISE(30 degree oblique) views.
- - localize soft tissue swelling.
- - small avulsion fracture of the tip of lateral malleolus.

X- ray examination is called for :

I. Pain around the malleolus
II. Inability to take weight on the ankle immediately after the injury
III. Inability to take 4 steps in emergency department
IV. Bone tenderness at the posterior edge or tip of medial or lateral malleolus

MRI…. Acute lat .collateral injury may be seen by MRI

Treatment:

.Initial treatment consist of RICE

Rest , Ice , Compression & Elevation

Which continue to 1-3 wk depending on the severity of the injury .

Ice (cold compress) should be applied for 20 min every 2 hr

More recently has been extended to

PRICER

Protection :by crutches ,splint or brace

Rehabilitation : supported return to function.

NSAID used in acute phase

Topical (NSAI) gel or cream is useful

Operative treatment

If the ankle dose not start to settle with in 1-2 wk of starting RICE further evaluation is needed .

Persistent problems at 12wk after the injury despite the physiotherapy , may signal the need for operative treatment .

Arthroscopic repair or ligament substitution is now effective in many cases



Deltoid lig tear:

Usually associated with the fractures of the distal end of the fibula or tearing of the distal tibiofibular lig. Or both

X-ray:

Widening of the medial joint space in the mortise view (s.t talar tilt or diastesis of the tibiofibular space)

Treatment

Provided the medial joint space are completely reduced the lig. Will heal

If it not reduce may need to be explored in order to free the soft tissue that tapped in the joint

Fractures of the ankle pott's fracture

Fracture & fracture - dislocation of the ankle are common .

The most obvious injury is a fracture of one or both malleoli & invisible part of injury (rupture of one or more ligaments which is just as serious .

Mech. Of injury :

The foot anchored to the ground while the body lunges forwards.

The ankle twisted , the talus tilt & or rotate in the mortise causing low injury fracture of one or both malleoli with or without lig. Injury

If the malleoli push off ,,,,, fracture obliquely

If the malleoli pull off ,,,,, fracture transverse

The fracture pattern depend on:

1. position of the foot
2. Direction of the force at the moment of injury

Danis & Weber classification which focus on the fibular fracture & according to the direction of the force.

Type A: is transverse # of fibula below the tibiofibular syndesmosis ,s.t. associated with oblique or vertical # medial malleolus.(adduction force)



Type B: oblique # of the fibula at the level of syndesmosis. Often there is also avulsion injury to the medial malleolus or torn medial (deltoid) ligament.(ext. Rotation)



Type C : severe form above level of syndesmosis-----tibiofibular ligament injured and interosseos membrane torn.(abd. and ext. rotation)

Clinical features:

- seen in footballer ,skiers , climbers and older group include women with post menopausal osteoporosis.
- History of severe twisting injury followed by

Intense pain ,inability to stand , swollen & deformity ankle.(suggest more serious than simple sprain



The site of tenderness is very important(if both the medial & lateral sides are tender ,, double injury must be suspected.

x-ray:

Ap, Lat & mortise views.

(fibula fracture is best seen in the lateral view). While diastases may not be seen without mortise view



Treatment

Guiding principles are:

1) Don't delay.(bec. Of the swelling) if the injury is not dealt with within a few hours , definitive treatment may have to be deferred for several days while the leg is elevated so that the swelling subside

2) Treat the entire injury , not only the fractures which are visible on x-ray, ligaments are not (like widening of the talo-fibular space , asymmetry of talo-tibial space ,tilted talus)

3) Reduce accurately.in assessing of acurat reduction

I. Fibulla restore it's full legnth
II. Talus sit squarely in the mortise
III. Medial jt. Space restored (4mm)
IV. No tibiofibular diastesis in oblique X-ray

4) Check and maintain reduction

the first is to decide whether the injury is stable or unstable.

This depends on tibiofibular ligament if torn or avulsed.

treatment

For undisplaced fractures:

- Conservative treatment with splinting and for Type B below knee cast is applied with ankle in neutral position ,check X-ray after 2 wk to confirm the fracture & the cast can discarded after 6-8 weeks.
- Ankle & knee movement must encouraged by active exercises

For displaced fractures:

reduction of joint disruption is mandatory fixation of fracture (for malleolar fracture internal fixation by one or two screws)

- closed reduction by traction and reverse the force of mechanism of injury..
- for displaced Weber Type C fracture:

  -the fibular # is above syndesmosis

  - 1st step is to reduce the fibula,restoring the length and stabilize by using plate and screws.
- syndesmosis is stabilized by insertion tranverse screws across the fibula into tibia.

Complications:

A) Early:

- vascular injury:especially in severe fracture and dislocation.----immediate reduction is mandatory.
- Wound break down: especially in D.M,wound edges break down(necrosis).
- Infection.

B) Late

1) incomplete reduction.

2) Non union:especially for medial malleolus because flap of periostum intrupt between fracture sides.

3) Joint stiffness.

4) Algodystrophy:pain in the foot,swelling,difuse tenderness.

5) Osteoarthritis : because of malunion , incomplete reduction-secondary osteoarthritis.

A

B

Ruedi classification



Pilon fractures

Mechanism of injury: occur when large forces drives the talus upward against the tibial plafond.

- High energy force responsible for most pilon fractures , but some caused by low energy rotational forces .
- There is considerable damage to the articular cartilage and subchondral bone may be broken into several pieces.

Clinical features:

Little swelling initially-------changes to fracture blisters

The ankle may be deformed and even displaced

x-ray

comminuted fracture of the distal end of the tibia , extending into the ankle joint.

In severe injury CT scan is mandatory

Treatment :

control soft tissue swelling is priority

This achieved by elevation and calcaneal traction.

Or by applying external fixator across the ankle.

Once the skin has recovered ,open reduction and fixation by plate and screw may be possible.

Pilon fracture takes 12-16 weeks to heal.



Medscape®
www.medscape.com

Complications:

I.Wound problems

II.Deep infections

III.Mal-union

IV. non-union

Thank you

BENIGN BONE TUMORS

Presented by
Dr. Karrar Kareem
Orthopedic surgeon
FIBMS, CABMS (Ortho)

Overview

- Bone tumors are very diverse in morphology and biological potential (can be no big deal or rapidly fatal)
- MOST bone tumors are benign lesions
- Most benign lesions are seen <30 years of age
- A new bone tumor in the elderly is more likely to be malignant
- Pathological fracture can be the first sign of tumor
- No bone is immune (though most primaries are in long bones)
- Location in the bone gives important Dx info
- More common benign lesions typically present as incidental findings (non-painful, stable size)

Classification of Benign Bone Tumors

Bony origin

i. Bone island

ii. Osteoma

iii. Osteoid osteoma

iv. osteoblastoma

Cartilaginous

i. Chondroma

ii. Chondroblastoma

iii. Chondromyxoid fibroma

iv. Osteochondroma

Fibrous

i. Fibrous cortical defect

ii. Non-ossifying fibroma (NOF)

iii. Desmoplastic fibroma

iv. Fibromatosis

Giant cell containing

i. Giant cell tumor (GCT)

ii. Aneurysmal bone cyst (ABC)

Classification of Benign Bone Tumors

Blood vessels

I. Hemangioma

II. Cystic angiomatosis

III. Hemangiomatosis

IV. Glomus tumor

Nerves

I. Neurofibroma

II.Neurilemmoma/Schwannoma

Fat

I. Lipoma

Epithelium

I. Implantation dermoid

Intraosseous ganglion

ii. Pigmented villonodular synovitis (PVNS)

iii. Synovial chondromatosis

iv. Lipoma arborescens

No known origin

i. Solitary bone cyst (SBC)

Non-neoplastic tumors

i. Brodie's abscess

ii. Hydatid

iii. Hematoma

iv. Infarction

v. Histiocytosis

Lytic benign bone tumors

- Osteoid osteoma
- Osteoblastoma
- Chondroma
- Chondroblastoma
- Chondromyxoid fibroma
- Fibrous cortical defect
- Non-ossifying fibroma
- GCT
- ABC
- Hemangioma
- Intraosseous lipoma
- SBC

Approaching a benign bone lesion

Clinical feature

May be less useful

As many lesions present with non-specific features of pain, swelling or pathological fracture.

Sex incidence is of little diagnostic value in primary bone tumors.

Plain radiograph

Mainstay of imaging modality

Views Antero-posterior, lateral and oblique

CT Scan

Can define matrix calcification

Fractures

Soft tissue extension

Approaching a benign bone lesion

MRI

Gives idea about varieties of tissues present

To access the extent of disease

Intraosseous edema and extraosseous inflammatory response may be demonstrated

Pre-op planning / post-op evaluation

Angiography

To see tumor vessels

To see encasement of major adjacent vessels

To distinguish between tumors …

e.g. Osteosarcoma is highly vascular and chondrosarcoma is hypovascular

Approaching…

Bone scintigraphy

Compound – technetium labelled diphosphonate

Phases

-- Blood pool phase

- Vascular lesion e.g. ABC ,
- Avascular lesion e.g. Cartilage tumors

-- Delayed phase

Helps to identify multiplicity of lesions

To see distant metastases

Patient is symptomatic (e.g. bone pain), but no lesion is identified in plain radiograph or CT Scan, then Bone scintigraphy can be helpful.

Technetium 99 bone scan

Scintigraphic study that makes use of Technetium99m (Tc99m-methylene diphosphonate (MDP)) as active agent.

3 stages which follow IV injection of the tracer.

- 1) Flow phase >> 2 to 5-sec images are obtained for 60 seconds after injection >> demonstrates perfusion and characterises the blood flow to a particular area
- 2) Blood pool phase >> the blood-pool image is obtained 5 min after injection >> demonstrated the blood pool, not the blood flow inflammation causes capillary dilatation and increased blood flow If the study is going to be a triphasic bone scan, a third phase is added.
- 3) Delayed phase >> the bone image is obtained 2 - 4 hours later

urinary excretion has decreased the amount of the radionuclide in soft tissue

BIOPSY

Most conclusive test because it confirms if the tumor is malignant or benign, the bone cancer type (primary or secondary bone cancer), and stage.

TYPES-

1. Needle biopsy: During this procedure, a small hole is made in the affected bone and a tissue sample from the tumor is removed. two types –

- Fine needle aspiration: During this procedure, the tissue sample is removed with a thin needle attached to a syringe.
- Core needle aspiration: During this procedure, the surgeon removes a small cylinder of tissue sample from the tumor with a rotating knife like device.

2. Incisional biopsy: During this procedure, the surgeon cuts into the tumor and removes a tissue sample

Fine and core needle

Many 'lumps and bumps' are diagnosed via fine needle aspiration which involves removing cells directly from the lump.





Core needle biopsy

True cut



Arch Bronconeumol. 2014;50:313–7

ENNEKING STAGING

Enneking described the most widely used staging system for benign bone tumors .

The stages are denoted by the Arabic numerals 1, 2, and 3

Stage 1-LATENT- low biological activity, well marginated , often incidental ,may resolve spontaneously. -NOF

Stage 2-ACTIVE-Symptomatic,limited bone destruction, may present with pathological fracture- ABC

Stage 3-AGGRESSIVE- Bone destruction/Soft tissue extension, require complete work-up and a removal with wide margins to avoid possible local recurrence. -GCT

Keys to characterize a lesion

Age
Location
Multiplicity
Shape
Size
X-ray appearance – lytic/dense
Zone of transition
Sclerosis
Periosteal reaction
Matrix calcification
Soft tissue involvement

Benign bone lesions

I. Bone island
II. Osteoma
III. Osteoid osteoma
IV. osteoblastoma

Bone island / Enostosis

Well demarcated round osteosclerotic in medulla of femur

Common

Usually incidental finding

Constitute a small focus of compact bone within a cancellous bone

One of the " don't touch lesions"

Age

Any age

May grow in size up to the age of skeletal maturity

Location

Can occur anywhere in the skeleton

some predilection for pelvis, long bones, spine and ribs.

Plain radiographic findings

Single or multiple

Round or oval

Always medullary in location

Has radiating thorn-like spicules with narrow zone of transition

Size – usually < 15mm , can be large up to 4 cm

No periosteal reaction, no cortical expansion

In diaphyseal bone, the long axis of a bone island typically parallels the long axis of the involved bone.

Bone island / Enostosis



Osteoma

skull

Seen almost exclusively in bones formed in membrane

Slow growing, usually asymptomatic, found incidentally

Location

Paranasal sinus

Skull vault

Mandible

Varies, some > 2.5 cm

Ivory/dense

(Lacks haversian system)

Spongy

(Resembles normal bone)

Radiographic features

Ivory osteomas appear very radio-dense

Spongy osteomas may demonstrate central marrow

Bone Scan >> Increased activity reflects the size of the lesion.

Complications

Growth within PNS can cause mucocele

Growth from inner table of skull may produce raised intracranial pressure

Association "Gardner syndrome"

> Multiple osteomas of skull, mandible and long bones

> Familial adenomatous polyposis

> Soft-tissue tumors of connective tissue origin



اصغر من ٢ سم + الم بالليل + يروح من ياخذ NSAID

Osteoid osteoma

10-35 yrs.

M:F = 3:1

Typical history of localized, intermittent bone pain of several weeks or months, occurring specially at night, with dramatic relief by NSAID's

Location

Long bones of limbs, majority in femur & tibia

Can occur anywhere

In spine, occurs in posterior elements.

(careful scrutiny of the neural arches at the apex of the concavity of scoliosis)

Pathology

An osteoid osteoma is composed of three concentric parts-

I. nidus

I. meshwork of dilated vessels, osteoblasts, osteoid and woven bone

II. may have a central region of mineralization

II. fibrovascular rim

III. surrounding reactive sclerosis

***The nidus releases prostaglandins (via Cox-1 and Cox-2) which in turn result in pain

Imaging

Plain radiograph & CT feature

>> A round or oval radiolucency with a sclerotic margin and contains a small dense center, known as nidus

>> Solid periosteal reaction with cortical thickening

>> Usually < 2 cm

>> Can be multifocal

Osteoid osteoma

Differentials

- Osteoblastoma (>2cm) does not respond to NSAIDs
- Stress fracture – that causes chronic cortical thickening
- Chronic osteomyelitis Sequestra

Area of radiolucency more irregular

Sequestra are irregular or linear in shape, differentiating from central nidus of osteoid osteoma

Osteoblastoma

Same as osteoid osteoma, but >2cm and no NSAID

Similar in histological characteristics as osteoid osteoma but usually larger

Less common than osteoid osteoma

80% of patients under the age of 30

In 2nd and 3rd decades

Clinically >> long insidious history of pain, often worsening at night

Not relieved by NSAID's

Limitation of movement

Location

Spine (posterior element), flat bones

Metaphysis and distal diaphysis of long bones.

Pathology

Larger than osteoid osteoma

Irregular in shape, friable and hemorrhagic

Abundant osteoid tissue and many thin-walled capillaries

Plain radiograph and CT features

2-10cm radiolucent lesion

Margin – irregular but sharply demarcated

Surrounding sclerosis – varies but can be profound

Can cause cortical expansion and thinning

Calcification or ossification of osteoid tissue – punctate or amorphous

Sometimes, lesion can be aggressive with soft tissue masses containing calcification and ossification

Bone scan

Active both in blood-pool and delayed phase

Chondroma/Enchondroma

=short bones, if long bone then calcium

2nd most common benign cartilage lesion (Osteochondroma is most common)

Single/multifocal

Usually central medullary

Childhood to adulthood, peak incidence = 20-50 years

Clinical feature

Rarely symptomatic, Patient may admit localized hard swelling for many years

>> Pathological fracture

Location

60% in hands – commonly in phalanges, less commonly in metacarpals

30% - long bones

10% - flat bones – pelvis, scapula and vertebral bodies.

Plain radiograph & CT scan

Cartilaginous tissue is not radiopaque

Radiolucent lesion

Expansion and thinning of overlying cortex

No periosteal reaction in the absence of fracture

Zone of transition is narrow and sclerotic

Endosteal scalloping = +- Flecks of calcification within the lesion, usually purely lytic in phalanges, in other locations, enchondromas are expansile, with characteristic "rings and arcs" calcifications.

MRI

T1 >> Low to intermediate signal

T2 >> typically of background intense high signal (high water content of hyaline cartilage matrix)

they can be focal regions of signal drop out where calcification present

Post-contrast

Variable enhancement

“rings and arcs” calcifications

purely lytic in phalanges

R

MRI & CT scan



OLLIER"S DISEASE

30% of malignancy

Ollier disease also known as enchondromatosis, is a non- hereditary, sporadic, skeletal disorder characterised by multiple enchondromas that are principally located in the metaphyseal regions.

Plain films show multiple enchondromas. Larger lesions can show cartilage calcification in a typical rings and arcs pattern.

Imaging characterestics are of same as ENCHONDROMAS

Risk of malignant transformation <30%



MAFUCCI'S SYNDROME

100% of malignancy

Maffucci syndrome is a congenital non hereditary mesodermal dysplasia characterised by multiple enchondromas with soft-tissue cavernous haemangiomas.

Imaging findings are multiple enchondromas seen associated with soft tissue swelling and phleboliths.

Overall risk of developing any malignancy up to 100%

Chondroblastoma

=epiphysis

Arises in the epiphyses or apophyses of long bones

Presents during the teenage years

The most common sites are the epiphysis of the proximal humerus, distal femur, and proximal tibia

Low-grade joint pain (constant, unrelated to activity) and swelling

Radiographic findings:

Small, well-defined lesions with a sclerotic border that may cross the physis

Matrix calcification may be seen

Chondroblastoma



Differential diagnosis

- Giant cell tumor
- Benign but locally aggressive skeletal tumor that occurs near the growth plate in young adults
- Chondromyxoid fibroma
- Avascular necrosis
- Clear cell chondrosarcoma
- Abnormality of subchondral bone in which pain is activity related. In contrast, in chondroblastoma, subchondral bone is normal, and pain is constant, unrelated to activity
- Aneurysmal bone cyst
- Osteomyelitis

NON OSSIFYING FIBROMA

Eccentric well-defined lytic lesion with sclerotic lobulated margin.

Also called Fibrous cortical defect or Metaphyseal cortical defect or Fibrous xanthomas

Musculoskeletal tumor - Occurs in 30 % of children - 1st two decades - Site : femur , tibia , humerus (8 % multiple lesions) - Asymptomatic

X ray : well defined , eccentric , radiolucent lesion in metaphysis ,multilocular app or rim of sclerosis , doesn't expand the cortex & no periosteal reaction.

Treatment : observation

NOF radiology

FIBROUS DYSPLASIA

Intrinsic defect of endochondral bone maturation immature ossification pattern

Characterized by replacement of normal bone & marrow by fibrous tissue & small woven spicules of bone weakbone .

Solitary or multifocal

- ALBRIGHT ' S SYN : (polyostotic F.D + café au lait spots + endocrinopathies .)

Site : rib , femur , humerus , tibia , maxilla

C.F : monostotic – asymptomatic bone pain , skeletal deformities

RADIOGRAPHIC APPEARENCE

Epiphysis or metaphysis or diaphysis

Well defined geographic lytic lesions

Ground glass matrix : intramedullary radiolucencies - Cortical thinning - Secondary deformities esp. in wt , bearing bones (microfractures – on tension side of bone)

Poorly defined areas of osteolysis suspect

Shepherd's crook deformity

Coxa varus angulation of the proximal femur, classically seen in femoral involvement by fibrous dysplasia, although may be seen in other disorders such as Paget disese of bone and osteogenesis imperfecta.

- The shape of the proximal femur resembles that of the staff carried by herders (shepherds), which is known as a crook

FIBROUS DYSPLASIA

Reumatol Clin. 2014;10:413–5

TREATMENT

Monostotic lesions, asymp

no Rx Indications

severe deformity

persistent pain

pathological #

- Curettage & grafting (cortical allograft prfrd. d/t slower absorption)

- Internal fixation +/ osteotomy

- Bisphosphonates

UNICAMERAL BONE CYST

90% found in the proximal part of the humerus and femur

Developmental anamoly of physis

Transient failure of ossification of physeal cartilage & cyst formation

< 20 yrs age , M > F -
Spontaneously resolve in late adolescence , rarely persist into adulthood

Site : proximal humerus , proximal femur , calcaneum .
Active cysts are juxtaposed to physis (within 1 cm)

C.F : usually asymptomatic , pathological #

Cysts progress from active to quiescent to involutional stage .

cysts usually shrink as patient approaches skeletal maturity.

Radiographic features : metaphysis of immature skeleton , DIGNOSTIC- well marginated , centrally located ,purely lytic lesion. radiolucent that expand and thin the cortex

FALLEN FRAGMENT SIGN – # fragment in the cyst



DD : ABC & fibrous dysplasia

Radiology

MRI & CT

MRI signal characteristics for an uncomplicated lesion include

T1 - low signal

T2 - high signal Usually there no fluid-fluid levels unless there has been a complication with haemorrhage

CT and MRI add little to the diagnosis, but are however helpful in eliminating other entities that can potentially mimic a simple bone cyst

TREATMENT

Small asymp lesions in upper extremities- observation & follow up Larger lesions

Symptomatic lesions need Rx

Lesions in lower extremities Various options are Curettage +/ - bone grafting +/- int. fixation

Aspiration & inj of steroids or bone products

Aneurysmal bone cyst (ABC)

=metaphysis of tibia and femur fluid-fluid levels

- Benign , Expansile
- Primarily seen in children and adolescents (80% l <20yrs of age)
- Blood-filled spaces of variable size separated by connective tissue (trabeculae of bone or osteoid tissue) and osteoclast giant cells
- Not lined by endothelium.

Types

- Primary
- Secondary (chondroblastoma, fibrous dysplasia, giant cell tumour (GCT), osteosarcoma)

Location and Imaging's

Long bones – tibia & fibula (24%), femur (13%),

Spine 20-30% (posterior elements). sacrum

- A variant of ABCs is the giant cell reparative granuloma.(SOLID ABC)
- seen in the tubular bones – hand , feet craniofacial skeleton.

ABC

BONE SCAN • Doughnut sign - increased uptake peripherally with a photopenic centre

X ray sharply defined, expansile, bubbly, osteolytic lesions, with thin sclerotic margins. Eccentricity is typical.

But very often missed out due to cortical thinning due to ballooning.

CT Demonstrates these findings to a greater degree, and is also better at assessing cortical breach and extension into soft tissues.

Fluid fluid levels (better than MRI)

Imaging



TREATMENT

Excision – cortical based or surface tumors

Extended intralesional curettage & grafting – central lesions

Embolisation – vertebral & pelvic ABC

Low dose radiation – effective method but not used d/t malignant transformation

Giant cell tumor

osteoclastoma

Knee, sub-articular

Distinct neoplasm arising from non-bone forming supportive connective tissue of marrow with network of stromal cells regularly interspersed with giant cells.

(Jaffe & Liechtenstein)

-75-80% of patients 20-50 yrs

Male:Female

-1:1.3 (Benign)

-3:1 (Malignant)

Giant cell tumor

55% AROUND THE KNEE

- 10% in the distal radius
- 6% in the proximal humerus
- SPINE rarely involved (commoner in the sacrum)
- In the head and neck region the maxilla and mandible are more commonly involved.

- SIGNS&SYMPTOMS

1. PAIN
2. SWELLING
3. JOINT RESTRICTION
4. MUSCLE WASTING
5. NEUROLOGICAL SIGNS
6. PATHOLOGICAL #
7. PULMONARY METS- 3 %

Pathology

GROSS

End of bone is expanded.

Eccentric lesion at the epiphyseo-metaphyseal region.

Thin periosteum.

Fleshy dark brown, soft, friable mass.

Cystic spaces seen

Microscopy

Vascularized network of round,oval or spindle shaped stromal cells and multinucleated giant cells(40-60 nuclei /cell) with numerous centrally placed nuclei



Grading system

Jaffe,Lichenstein and Portis(1940)

GRADE 1- Conventional GCT

GRADE 2- Boderline tumours

GRADE 3- Sarcomatous type of stroma



Radiology

Trabaculation pure lytic (60%)

Fine trabacultion(40%)



Imaging



Imaging

Surgical Treatment

Stage1& Stage2 --- Intralesional or Marginal Excision

Stage3 --- Wide resection with Reconstruction

Radiation, Embolaisation



Curettage & bone grafting indication stage-1&2

Adequate window

Motorised burr

Bisphosphonates- locally or systemically prevent recurrence.

a
b
c
d
e

B
C

Thank you

Malignant bone tumor

Dr. Karrar Kareem

Double board certified

Malignant bone tumors

Can be classified as

Primary (arising from abnormal bone or cartilage Cells)

Secondary (bone metastases of other tumors).

The most common primary bone tumors are

- Osteosarcomas
- Ewing sarcomas
- Chondrosarcomas

Progressive, localized pain that worsens at night or with physical activity is common and usually accompanied by swelling

Malignant bone tumors…

In Ewing sarcoma and osteosarcoma, symptoms often first manifest after an injury.

Diagnosis is confirmed by imaging and biopsy

Depending on the type and stage of the tumor

Osteosarcomas

chemotherapy, radiation therapy, and/or definitive surgical resection may be required.

Bone tumors may also occur secondary to metastases

of other primary cancers.

The metastases are commonly detected in the spine and pelvis and usually arise from the lung, breast, and prostate cancer

Musculoskeletal Tumor Society

MSTS Staging for Malignant tumors

Stage	Grade	Site	Metastasis
IA	Low Grade	T1 - intracompartmental	M0 (none)
IB	Low Grade	T2 - extracompartmental	M0 (none)
IIA	High Grade	T1 - intracompartmenta	M0 (none)
IIB	High Grade	T2 - extracompartmental	M0 (none)
III	Metastatic	T1 or T2 - intra or extra-compartmental	M1 (regional or distant)

American Joint Committee on Cancer

AJCC Staging System

Stage	Grade	Size	Depth	Node	Metastasis	5 yr. survival
IA	Low Grade	< 8cm	Any	None	None	98%
IB	Low Grade	> 8cm	Any	None	None	
IIA	High Grade	< 8cm	Any	None	None	82%
IIB	High Grade	> 8cm	Superficial	None	None	82%
III	Any	Discontinuous (skip) lesion	Deep	None	None	52%
IVA	Any	Any	Any	None	Lung	
IVB	Any	Any	Any	Present	Other than lung	30%

Compartments and Grade

Tumor Compartments

Intracompartmental

bone tumors are confined within the cortex of the bone

Extracompartmental

bone tumors extend beyond the bone cortex

Tumor Grade

Histologically, tumors are graded based on the percentage of cellular atypia

Low grade tumors

low metastatic potential

e.g. parosteal osteosarcoma

High grade tumors

greater metastastatic potential

e.g. intramedullary osteosarcoma, Ewing's sarcoma, dedifferentiated chondrosarcoma

Prognosis

Prognosis of bone cancer depends on the following variables

- Overall stage of disease (incorporates all of the below)
- Presence of metastasis
- Skip (discontinous) lesions within the same bone
- Histologic grade
- Tumor size

Ewing sarcoma

Definition

Highly malignant bone tumor arising from neuroectodermal cells

Some sources suggest that Ewing sarcoma originates from mesenchymal stem cells

Etiology:

Associated with various chromosomal translocations of the EWSR1 gene (chromosome 22)

EW =Ewing
SR=sarcomas

t(11;22)

Incidence

Peak at 10–20 years

Kids

Sex: ♂ > ♀

Ethnicity: primarily affects white individuals

Ewing sarcoma

Primary tumor: often diaphysis of long bones (particularly femur, tibia, fibula, and humerus) and bones of the pelvis

Metastasis

lungs, skeletal system, and bone marrow

Clinical features

localized pain (progressive, worsens at night), hyperthermia, and swelling after trauma to the bone (tissue mass that is tender to palpation; and accompanied by erythema)

Diagnostics

Conventional X-ray

Lytic bone lesions

Onion skin appearance of the periosteum



Biopsy

Anaplastic small-blue-round cell malignancy

Tumor cells resemble lymphocytes

Differential diagnoses include

lymphoma

chronic osteomyelitis

Chromosomal translocation t(11;22) which leads to expression of fusion protein EWS-FLI1

most common translocation diagnostic of Ewing sarcoma and is present in more than 90% of cases.

Cells contain glycogen accumulations and are usually CD99-positive

Laboratory findings:

↑ ESR

↑ LDH

leukocytosis

"Ew, did you feed on 22 onions?": Ewing sarcoma, femur region, chromosome 22, onion skin appearance.

Imaging's X Ray



Ewing Sarcoma MRI



Treatment

Surgery; (definitive resection) plus neo-adjuvant and adjuvant poly-chemotherapy

Includes a combination of vincristine, doxorubicin, ifosfamide, and etoposide.

Additionally: radiation therapy

If complete resection is uncertain or impossible,
if chemotherapy is insufficient,
metastases

Prognosis

Extremely aggressive, early metastases

Usually responsive to chemotherapy

Five-year survival rate of ~ 80% for localized disease

Osteosarcoma

Malignant, osteoid

, and bone-forming tumor arising from mesenchymal stem cells (osteoblasts) located in the periosteum

Etiology

Primary osteosarcoma: unknown

Secondary osteosarcoma

Paget disease of bone

Rdiation injury

Bone infarction

1000 times!!!!

Increased incidence in individuals with retinoblastoma and Li-Fraumeni syndrome

Osteosarcoma

Epidemiology

Bimodal distribution

Primary osteosarcoma

in puberty/adolescence age 10-30 years

Secondary osteosarcoma advanced age

Sex: ♂ > ♀

Most common primary malignant bone tumor

Localization

Primary tumor **metaphyses** of long bones (particularly distal femur and proximal tibia)

Metastases

lungs, skeletal system, regional lymph nodes

Most common primary bone malignancy, but Multiple myeloma, which is sometimes considered a primary bone tumor, is more common than osteosarcoma.

Clinical features

Frequently first manifests with pain (progressive, worsens at night and with activity)

Progressive swelling (tissue mass that is tender to palpation; and accompanied by erythema)

Pathologic fractures

Limping, decreased range of motion

Possible benign symptoms

Imaging

Conventional x-ray

Sunburst appearance of lytic bone lesions and/or **Codman triangles**

Signs of osteolysis adjacent to osteosclerosis (moth-eaten appearance)

MRI

assesses the involvement of soft tissue

, evaluation in cases of unclear radiographic findings

Osteosarcoma



Investigations

Biopsy

Pleomorphic, malignant osteoblasts that produce osteoid

Osteosarcomas always feature woven bone matrix (compared to chondrosarcomas and fibrosarcomas)

Laboratory

↑ Alkaline phosphatase

↑ LDH

↑ ESR

Treatment

Surgery; (definitive resection) with neoadjuvant and adjuvant polychemotherapy (e.g., a combination of methotrexate , doxorubicin, cisplatin, and ifosfamide)

Osteosarcomas are usually **resistant to radiation therapy**

Histological examination of the resected bone to evaluate the effect of neoadjuvant chemotherapy

(major prognostic factor)

Prognosis

Aggressive course

Primary osteosarcoma five-year survival rate of ~ 70% (usually responsive to treatment)

poor prognosis

Secondary osteosarcoma

: poor prognosis (less responsive to treatment)

Chondrosarcoma

Definition: a malignant tumor arising from mesenchymal cells that produce cartilage

Etiology

Primary chondrosarcoma: unknown

Secondary chondrosarcoma: e.g., osteochondroma, Paget disease of bone, radiation

Epidemiology

Age: usually > 50 years

Sex: ♂ > ♀

Localization: most common in the medullary cavity of the pelvis, ribs, proximal femur, and proximal humerus

Clinical features

Deep, dull pain (worsens at night, insidious progression over months to years)

Local swelling

Pathological fractures

Neurovascular disturbances and/or limited range of motion

Diagnostics

Conventional X-ray or CT

Osteolysis with a **moth-eaten appearance**

Calcifications (rings and arcs Enchondroma calcification, popcorn calcification)

Endosteal scalloping and cortical breach with infiltration of soft tissue

MRI rim-like contrast enhancement



Chondrosarcoma





RT

Biopsy

Malignant chondrocytes

Lobulated appearance (hyaline cartilage nodules with peripheral calcification)

Grading

- Grade I: low cellularity, mostly chondroid matrix
- Grade II: increased cellularity, decreased chondroid matrix with localized myxoid changes
- Grade III: high cellularity, prominent nuclear atypia , myxoid matrix

Treatment

Surgery (definitive resection)

± Chemotherapy and radiation therapy

, possibly as adjuvant therapy or as palliative treatment

Prognosis

Five-year survival rate of 50–85% (depending on the histological grading)

Late recurrences are possible.

Regular follow-ups for 10 years are required.

Chordoma

Chordomas are malignant tumors of primitive notochord origin that most commonly occur in the sacrum and coccyx.

The tumor typically presents in patients > 50 years of age with insidious low back pain with bowel or bladder dysfunction

Epidemiology

Most common primary malignant spinal tumor in adults

3:1 male to female ratio

Usually in patients > 50 years

50% occur in the sacrum and coccyx

35% in spheno-occiptal region

15% in mobile spine

Presentation

Pain

Insidious onset of pain

may be mistaken for low back or hip pain

Neurologic

often complain of bowel or bladder changes

sensory deficits rare due to distal nature of tumor

Gastrointestinal

constipation

fecal incontinence

Physical exam

Neurologic

motor deficits rare because most lesions at S1 or distal _

bowel and bladder changes are common

Rectal exam

more than 50% of sacral chordomas are palpable on rectal exam

Imaging

Radiographs

often difficult to see lesion due to overlying bowel gas

CT

will show midline bone destruction and soft tissue mass

calcifications often present within the soft tissue lesion

MRI

bright on T2

useful to evaluate soft tissue extension



Treatment

Nonoperative

radiation treatment

inoperable tumors

Operative

wide margin surgical resection +/- radiation

standard of care in most patients

technique

must be willing to sacrifice sacral nerve roots to obtain adequate surgical margins

… Add radiation if margin not achieved

Secondary malignancies of the bone (bone metastasis)

secondary bone tumors due to metastasis

(predominantly hematogenous) of primary malignancies of other organs

Etiology: : primary malignancies (most commonly lung, breast, and prostate cancer

, together comprising ~ 80% of cases)

Epidemiology

More common than primary bone tumors (~ 70% of all malignant bone tumors)

Bones are the third most common site of metastases, after the lung and the liver

Localization: most commonly spine (~ 40%) and pelvis (~ 20%)

Classification

Osteoblastic metastasis

New bone formation
by osteoblasts outweighs osteolytic processes →
increase in radiographic density

Examples: prostate cancer, small cell lung cancer

Osteolytic metastasis

Osteolytic processes outweigh new bone formation
→ decrease in radiographic density

Examples: multiple myeloma, thyroid cancer, kidney cancer, melanoma, non-small cell lung cancer

Mixed metastasis

e.g., breast cancer, gastrointestinal cancer

Clinical features

Often no or few symptoms (incidental finding during routine staging of the primary tumor)

Local pain

Swelling

Pathologic fractures

Spinal cord compression

, and/or radicular symptoms

Approach

Imaging is generally only performed in patients with primary malignancies who have clinical signs or laboratory findings (e.g., elevated alkaline phosphatase, hypercalcemia) indicating metastatic bone disease

Biopsy

is performed to confirm the diagnosis when imaging is insufficient or the patient has no known history of cancer.

Laboratory

Elevated serum calcium

Radiographic imaging

Choice of imaging studies depends on:

- The location of the suspected lesion and the clinical presentation
- The most likely underlying primary tumor

For suspected spinal lesions:

- Contrast-enhanced MRI (first-line modality): used to detect tumor extension and possible spinal cord compression
- CT may be performed to assess vertebral bone integrity.

For suspected extremity lesions:

- Conventional X-ray (initial test)
- Followed by CT or MRI if a pathological fracture is identified or suspected for operative planning

Whole-body skeletal imaging: indicated if a metastatic lesion is identified to detect additional, asymptomatic sites of metastasis

Radiographic imaging

Bone scintigraphy (bone scan): most commonly used test to detect metastases within the entire skeleton and indicate biological activity of lesions (osteoblastic, osteolytic)

PET scan: useful to detect metastases

and for staging, particularly for bone metastases

from breast and lung cancer

X-ray skeletal survey: primarily used for purely lytic metastatic bone disease (particularly multiple myeloma)

Whole-body MRI or CT are less sensitive than bone scintigraphy or PET/CT.

Biopsy

Preferably taken from a soft tissue mass at the afflicted site, otherwise from an accessible bone where biopsy does not cause lasting impairment of motility or stability

In the case of an isolated lesion, it should be presumed to be a primary tumor unless proven otherwise

Treatment

The choice of treatment generally depends on the underlying primary malignancy and stage of the disease.

Pain management

Systemic therapy

Chemotherapy

Osteoclast inhibitors (e.g., bisphosphonates): inhibit bone resorption

Localized tumor radiation

Surgery

If pathologic fractures have occurred or are likely: stabilize and restore function

If spinal instability or spinal cord compression has occurred or is likely: immediate decompression and stabilization

A 15-year old male without previous medical history has been complaining of right shoulder pain for 1 month. He also reports that he has lost 5 pounds within that same timeframe. He states that he has difficulty sleeping at night due to feeling warm and sweaty. Current imaging is shown in Figures A and B. A bone marrow biopsy is performed and is found to be positive for a EWS-FLI1 fusion protein. Which of the following combines the correct diagnosis with the correct translocation?



1. Synovial sarcoma; t(x;18)
2. Ewing's sarcoma; t(12;22)
3. Osteosarcoma; t(9;22)
4. Osteosarcoma; t(x;18)
5. Ewing's sarcoma; t(11;22)

THANK YOU

Spine and spinal cord trauma

Presented by

Dr. Karrar Kareem

CABMS, FICMS (ORTHO)

- Annually 15,000/year permanent spinal cord injuries in USA
- Commonly men 15-30 years old
- Nearly 10,000 peoples die each year due to spinal cord injury
- Male sustain SCI 3-4 time more than females.
- 15% patient develops neurological deficit.
- Mechanism of Injury RTAs: 50% Falls: 30% Penetrating Trauma: 10% Sport Injuries: 10%

- 10% of all spinal cord injuries occur from improper handling of the spine and patient after injury
- ASSUME based upon MOI that patient's have a spinal injury
- Manage ALL spinal injuries with immediate and continued care
- Lifelong care for spinal cord injury victim exceeds $1 million.
- Best form of care is public safety and prevention programs.

INCIDENCE SPINE INJURIES

Spinal segment and level of injury

Incidence

- Cervical 60%
- Thoracic 8%
- Thoracolumbar 20%
- Lumbar 10%
- Sacral 2%

T11-L2 because this site is transition between the rigid thoracic (dorsal) to the flexible lumbar spine.

Cervical spine injury and Neurological Deficit

Incidence of Neurological Deficit

1. Atlanto -occipital dislocation Up to 100%
2. Atlas 1-2% Axis 10% C3-T1 6%
3. Unilateral cervical facet dislocation 60%
4. Bilateral c. facet dislocation Up to 100%

Spinal Anatomy and Physiology

The spinal cord extends from the midbrain at the base of the skull to the level of L1 or L2 in most adults

31 pairs of peripheral nerves (spinal nerves)

8 cervical, 12 Thoracic, 5 Lumbar, 5 Sacral, and 1 Coccygeal Spinal Anatomy


anterior view
right lateral view
posterior view
cervical vertebrae (C1-C7)
thoracic vertebrae (T1-T12)
lumbar vertebrae (L1-L5)
Atlas (C1)
Axis (C2)
C7
T1
T12
L1
L5
sacrum (S1-S5)
coccyx
© 2013 Encyclopædia Britannica, Inc.
The structure of the segments of the spine
Cervical
Thoracic
Lumbar
Sacrum
Coccyx
Transverse process
Vertebral disc
Body
Spinous process
Vertebral foramen
Lamina
Articular process



a The superficial anatomy and orientation of the adult spinal cord. The numbers to the left identify the spinal nerves and indicate where the nerve roots leave the vertebral canal. The adult spinal cord extends from the brain only to the level of vertebrae L_1–L_2; the spinal segments found at representative locations are indicated in the cross sections.

b Lateral view of adult vertebrae and spinal cord. Note that the spinal cord segments for S_1–S_5 are level with the T_{12}–L_1 vertebrae.

c Inferior views of cross sections through representative segments of the spinal cord, showing the arrangement of gray matter and white matter.

Function of vertebral column

1.Supports the head.

2. Help maintain balance in the upright position.

3. Enclose and protect the spinal cord.

4. Permits movement (move forward, backward, sideways, and rotate).

5. Absorbs shocks during walking.

6. Serve as a point of attachment for the ribs, pelvic girdle and muscles of the back and upper limbs







Figure 6-4 & 6-5. (4) Cross-section of the cervical spinal cord. The diagram shows the anterior and posterior spinal arteries with their branches and territories. (5) Vascularization of the spinal cord: ventral vie. In: Waxman SG. *Clinical Neuroanatomy*. 26th ed. http://www.accessmedicine.com. Accessed November 2, 2009.


Posterior external venous plexus
Posterior internal venous plexus
Posterior intermediate spinal vein
Venous vasocorona
Posterior median spinal vein
Posterior radicular and spinal medullary veins
Dura mater
Anterior radicular and spinal medullary veins
Anterior median spinal vein
Anterolateral spinal vein
To anterior external venous plexus
Anterior internal venous plexus

Venous drainage

- The spinal veins form loose-knit plexuses in which there are an anterior and a posterior midline longitudinal vein, and on each side a pair of longitudinal veins posterior to
- the anterior and posterior nerve roots.
- These veins drain to the internal vertebral venous plexus (1) (between dural and bony canal) , and thence via the external vertebral venous plexus (2) to the segmental veins: vertebral in the neck; azygos in the thorax; lumbar in the lumbar region; and lateral sacral in the sacral region.
- At the foramen magnum they communicate with the veins of the medulla.



Division of vertebral column

C-spine

Including 7 vert.

Sole support of the head

- Head weighs. about 16-22 pounds 7-9 kg

Atlas- C1

Support the head

Securily affixed the occiput

Permitts nodding

It has no vert. Body or spinous process


SUPERIOR VIEW OF C1 ON C2
Axis: spinous process
Superior articular facet, atlas
Posterior arch, atlas
Dens
Transverse lig., atlas
Anterior arch, atlas
Jefferson Fracture
Lateral Mass Fx
Bony Avulsion TL
Midsub. Tear TL


The first two cervical vertebrae: the atlas and the axis
Anterior arch of atlas
Atlas
Dens (odontoid process)
Joint that permits nodding (as in indicating "yes")
Ligament that enables rotation (as in shaking the head to indicate "no")
Axis
Posterior arch of atlas

AXIS C2

- Odontoid process
- Projects upward
- Provides pivot point so the head can rotates

C7 VERTEBRA

- Prominent spinous process
- Vertebrl a. don't pass through its transverse foramina


Vertebral Arteries
Posterior cerebral artery
To Brain
Basilar artery
Skull
Intracranial, Intradural (V4)
Atlantal, Suboccipital (V3)
C1 (Atlas)
Transverse processes
C2 (Axis)
C3
Foraminal (V2)
C4
Vertebral artery
C5
C6
Preforaminal (V1)
C7
Ribs
Subclavian artery
T1
Aorta
THESE IMAGES ARE COPYRIGHTED BY AMICUS VISUAL SOLUTIONS.
COPYRIGHT LAW ALLOWS A $150,000 PENALTY FOR UNAUTHORIZED USE.
CALL 1-877-303-1952 FOR LICENSE.


Posterior cerebral arteries
Basilar artery
V4 (intradural)
V3 (C2 to dura)
V2 (foraminal)
V1 (pre-foraminal)

THORACIC SPINE

- 12 Ribs
- 1st rib articulate with the T1

  Attached to the transverse process and vertebral body

- Next nine ribs attached to the sup. and inf. Portion of adjacent vertebral bodies

  limits rib movement and provide rigid stability

- Larger and stronger than c- spine


Body
Vertebral foramen
Superior vertebral notch (forms lower margin of intervertebral foramen)
Superior costal facet
Pedicle
Angle of articular facet
Transverse costal facet
Lamina
Superior articular facet
Spinous process
Vertebral canal
Superior articular process and facet
7th rib
Spinous process (T 7)
Transverse process (T 9)
Lamina
Inferior articular process (T 9)
Spinous process (T 9)
Body
Transverse process
Costal facet
Inferior articular process and facet
12th thoracic vertebra: lateral view

LUMBAR SPINE

- 5 vertebrae
- Bear forces of bending and lifting above the pelvis
- Largest and thickest vertebral bodies and I.V disks


2nd lumbar vertebra:
superior view
Vertebral canal
Superior
articular process
Mamillary process
Transverse
process
Vertebral body
Accessory
process
Spinous
process
Lamina
Inferior
articular process
3rd and 4th lumbar vertebrae:
posterior view
Transverse process
Spinous process
Vertebral body
Intervertebral disc
Inferior
articular
process
Inferior
vertebral notch
Intervertebral
foramen
Superior
vertebral
notch
1
2
3
4
5
Articular facet
for sacrum
Lumbar vertebrae, assembled:
left lateral view

SACRAL SPINE

- 5 fused vertebrae
- Form posterior plate of the pelvis
- Protect the genitourinary systems
- Connect the pelvis and lower limbs to the axial skeleton



Figure 07.20ab Tortora - PHA 11/e

Copyright © John Wiley and Sons, Inc. All rights reserved.

Types of mechanism of spinal injuries



Spinal Column Injuries

Dislocations and subluxation

Fractures includes

- Spinous or transverse process
- Vertebral body
- Ruptured I.V disks

Common site of injuries

- C1-C2 delicate vert.
- C7
- T12-L1

Spinal Cord Injuries

- Concussion

  Similar to the cerebral concussion

  Temporary and transient dysfunction of cord

- Contusion

  Bruising of the cord

  Tissue damage, vascular leakage and swelling

- Compression

  Secondary to

  - Displacement of vertebra
  - Herniation of I.V disk
  - Displacement of vertebral bone fragment
  - Swelling from adjacent tissues

Cont…

- Laceration

Causes

- Bony fragment driven into the cord
- Stretching up to point of injury

- Hemorrhage


Hemicord Syndrome
Dorsal Column
(vibration + proprioception)
Ipsilateral
contralateral
Pain + temp
loss
Ipsilateral
motor
loss
Ipsilateral
vibration +
Proprioception
loss
Corticospinal
tract
(Motor)
Ipsilateral
Spinothalamic
Tract
(Pain + Temperature)
Contralateral


Descending Tracts (Motor)
Ascending Tracts (Sensory)
Lateral Corticospinal
Tract (Motor)
Doral Columns
(Deep touch, Propioception,
Vibration)
Lateral Spinothalamic Tract
(Pain, Temperture)
Ventral Corticospinal
Tract (Motor)
Ventral Spinothalamic Tract
(Light touch)

Spinal cord transection either complete or incomplete

Complete

No preservation on any motor or sensory function more than 3 segments below level of injury

C- spine >>> quadriplegia, incontinence, and resp. paralysis

Below T1 >>> incontinence, paraplegia

incomplete

INCOMPLETE
INJURY
Check for
sacral sparing!


Incomplete lesions of the spinal cord
Central Cord Syndrome
Anterior Cord Syndrome
Brown-Séquard Syndrome
Corticospinal tract
Spinothalamic tract

Incomplete>>> Anterior Cord Syndrome

- Crush injury or compression from hematoma
- Compression of anterior spinal artery
- Paraplegia below level of lesion
- Pain and temp. loss below the lesion
- Sparing of dorsal column transection


Summary of Spinal Cord Syndromes
Central Cord Syndrome
Incomplete loss of motor function affecting upper extremities greater than lower.
Anterior Cord Syndrome
Complete Motor Dysfunction below the level of the lesion with pain and temperature sensory loss, intact vibratory sensation and proprioception.
Brown-Sequard Syndrome
Corticospinal tract
Spinothalamic tract
Loss of motor function with vibratory and proprioception deficits on ipsilateral side of cord lesion.
Loss of pain and temperature sensation on contralateral side of lesion.

Incomplete>>> Central Cord Syndrome

- Most common type
- Better prognosis than Ant. Cord
- Hyperextension injury
- Bladder dysfunction
- More dense in the upper limbs and peripheral parts
- Sparing of light touch and proprioception


Dorsal columns
Lateral corticospinal tract
Lateral spinothalamic tract
Most commonly caused by **extension** injury
Upper extremities > **Lower** extremities
Distal muscles > **Proximal** muscles
Loss of motor function
Incomplete loss of motor function
Variable degree of sensory loss

Incomplete>>> Brown Sequard Syndrome

- Hemi-section of spinal cord
- Usually penetrating trauma
- **Best** prognosis
- **Rare** type
- Epsilateral motor, proprioception, light tough below the lesion
- Contralateral pain and temp. 1-2 segments below the lesion

Brown-Séquard Syndrome



Causes

Trauma (gunshot wound, stabbing)
Spinal cord tumor
Ischemia
Infection



Syndrome	Mechanism	Clinical	Prognosis
Anterior cord	Flexion or vascular	Complete loss of motor, pain, & temperature below injury, but retains proprioception and vibratory sensation	Poor
Central cord	Forced hyperextension	Sensory and motor deficit Upper > Lower Extremities	Average
Brown-Séquard	**Penetrating trauma**	**Ipsilateral loss of motor, vibratory sensation, and proprioception with contralateral loss of pain and temperature sensation**	**Good**

Ipsilateral loss of motor, vibratory, and proprioception

Contralateral loss of pain and temperature sensation

Fractures

>> 90% are simple fractures and it will heal with rest and symptomatic management

>>Unstable c- spine fractures need careful immobilization

…. These injuries should be reduced as soon as possible by any methods(traction or ACDF)

JEFFERSON #, ATLAS # C1

- C1 # usually follow axial loading injury or compression mechanism
- Can be assessed by open mouth view or CT scan
- If TAL(transverse atlantal lig.) is ruptured >> **unstable injury**

C1 'Jefferson' fracture - Open mouth view

Normal - open mouth view





treatment

- Halo jacket for 3 months is the usual treatment
- persistant instability require surgery in form of (post. Occipito-cervical fusion)

Hangman fracture(C-2)

- Fracture of C2 pedicle
- 5-10 % are caused by Hyperextension injury
- Very fatal injury



Odontoid fracture of C-2, Axis

- Most commonly missed c- spine injuries
- Constitutes about 10% of cervical injuries
- May lead to serious complication and death
- Rx mainly Halo jacket


Odontoid fracture classification
Type I :
Fracture of the upper part of the odontoid peg ; it's rare and potentially unstable
Type II :
Fracture at the base of the odontoid ; unstable, and has a high risk of non-union
Type III :
Through the odontoid and into the lateral masses of C2 ; best prognosis for healing



Flexion tear drop fracture

- Sudden hyper flexion and axial compression
- Constitutes 15% of c- spine injuries
- Involve disruption of all column
- Highly unstable
- Usually presented with neurological injury>>>ANT. CORD
- Rx mainly surgical by anterior plate stabilization





Bilteral facet dislocation

- Hyper extension with rotation(diving, MVA)
- Highly unstable
- Disruption of all spinal column ligaments
- Almost always Quadriplegic

>>>> poor prognosis



Spinal cord injury without radiographic abnormality(SCIWORA)

- Defined as traumatic injury with absence of finding on (plain x-ray, flexion-extension radiograph and CT scan)
- Usual mechanism is acceleration-deceleration or rotation injury
- If its suspected MRI should be done
- They require immobilization to prevent further injury



THORACOLUMBER REGION>> mechanism of injury

- Low-Energy Insufficiency Fractures arising in osteoporotic bone
- Minor Fractures of the Vertebral Processes due to compressive, tensile or torsional strains
- High-Energy Fractures or Fracture-Dislocations due to major injuries sustained in motor vehicle collisions, falls or diving from heights, sporting events, horse-riding and collapsed buildings.
- Neurological complications are mainly associated with the third group.

Mechanism …

- **Flexion Compression** – failure of the anterior column and wedge-compression of the vertebral body.
- **Lateral compression** – lateral wedging of the vertebral body resulting in a localized 'scoliotic' deformity.
- **Axial compression** – failure of anterior and middle columns >> Often unstable.
- **Flexion–rotation** – failure of all three columns and a risk of displacement or dislocation >> Usually unstable.
- **Flexion–distraction** – the so-called 'jack-knife' injury causing failure of the posterior and middle columns and sometimes also anterior compression >> Usually unstable.
- **Extension** – tensile failure of the anterior column and compression failure of the posterior column >> Unstable.

Classification systems

Denis
3-column classification



AO
classification



TLICS
classification



TLICS	3 independent predictors			
1	**Morphology** immediate stability	- Compression - Burst - Translation/rotation - Distraction	1 2 3 4	- Radiographs - CT
2	**Integrity of PLC** longterm stability	- Intact - Suspected - Injured	0 2 3	- MRI
3	**Neurological status**	- Intact - Nerve root - Complete cord - Incomplete cord - Cauda equina	0 2 2 3 3	- Physical examination
Predicts		- **Need for surgery**	0 – 3 4 > 4	- nonsurgical - surgeon's choice - surgical

COLUMN THEORY - DENIS

>> Anterior- ALL , anterior 2/3 body

>> Middle - post 1/3 body, PLL

>> Posterior- all structures posterior to PLL

Spinal injury and Three column concept:

- One column injury is **stable**
- Two column injury is >> **unstable**
- Three column injury is >> **invariably unstable**



Denis Classification of Spinal Trauma



Major Injuries

COMPRESSION
BURST
SEAT-BELT-TYPE
FRACTURE-DISLOCATION

Minor Injuries

transverse processes fx
articular process fx
pars interarticularis fx
spinous process fx

Compression (wedge) fracture

- 45% of all TL #s
- Anterior column failure
  (ant. Or lat. flexion)
- Middle , post. Column intact
- Usually no neurological deficit

Burst #s

- 15% of all TL #s
- Anterior and middle column failure

(Axial compression)

- Most common at the TL junction
- Ass. With neurological deficit



Flexion distraction=chance #=seat belt#

- Posterior and middle column failure (Hyper flexion then tension force)
- Anterior column act as hinge point and may damaged



Fracture dislocation

- Failure of all columns (compression, tension, rotation, or shear)
- Anterior hinge is disrupted
- Dislocation
- Severe neurological deficit
- Invariably unstable



General approach to spine trauma

Pre-hospital care

>>proper extraction and immobilization

…cervical collar

…sand bag

…hard board(log roll)

…tape

>>air way protection

>>rapid and safe transfer



Emergency assessment

- ABC and Immobilization

>>> hemodynamically stable

- Secondary survey

…log roll tech.

…remove spinal board

…remove c-spine collar carefully

…brief hx (mechanism, movement, position)

inspection

- Clothes removed
- Bleeding, abrasions , laceration
- Deformity, swelling
- Limb asymmetry and movement
- Chest expansion>>paradoxical movement

Palpation

- Tenderness
- Swelling
- Inter-spinous wedging > 7 mm
- Malalignment of spinous process
- Step-off

- **Must see pt's face – tenderness – how?**
- Identify important anatomical landmarks



Initial assessment





Sensory examination (dermatomes and reflexes

- Babiniski sign
- Perianal sensation
- Rectal tone
- Cremastric reflex



Radiological assessment

- X-ray
  - AP View
  - Lateral view



CT Scan

- **Assessment of**

>>> canal compromise

>>>comminution

>>>dislocation





MRI

Accurate assessment of soft tissues

- cord lesion
- ligament injury
- disc herniation
- hematoma

MANAGMENT

- Goals of Mx
- Maximize neurological recovery
- Maintain or restore spinal alignment
- Obtain healed and stable spinal column
- Prevent future deformity

- Medical treatment
- Methylprednisolone

>> 30 mg/kg bolus over 15 minutes

>>5.4mg/kg/h infusion over 23 hrs(1st 3hrs)

>> 5.4mg/kg/h infusion over 47 hrs(if more than 8 hrs passed)

- PPI
- LMWH

NON-SURGICAL Mx

- INDICATIONS
- **anterior wedging < 40%**
- **canal compromize < 40%**
- **kyphosis <25 degrees**

- **Rest**
- **Bracing**

Thank you

Introduction to Cardiothoracic Surgery

Assisted professor

Laith Sharba

MBChB FICMS EACTS FACS

Medical College – JIHMU

Consultant Cardiothoracic and Vascular surgeon

JMU University
Faculty of Medicine
Surgical department
Cardiothoracic & Vascular surgery

Course Title:
Cardiothoracic & Vascular surgery
Course Calendar: **Theory (14) hours**
Clinical (15) hours
Total credit:

Placement: **Fifth year 2023/2024**

Theoretical instructor:
Assistant Professor Dr. Laith Fathi Sharba
Consultant of Cardiothoracic & vascular surgery

Fifth Year`s Curriculum 2023-2024
Theoretical content

ت	أسم المحاضرة	عدد الساعات
1	Introduction to Thoracic Surgery, Anatomy, physiology & Symptomatology	
2	Diseases of the pleura	2
3	Diseases of the Lung	2
4	Neoplasm of the lung	2
5	Endoscopic thoracic surgery Part 1: Bronchoscope & esophagoscope	1
6	Part 2: Mediastinoscope	1
7	Part 3: Thoracoscope	1
8	Diseases of esophagus	3
9	Cardiac Surgery	2

Learning Resources:

Blackboard, Overhead Projector, Posters, Slideshow, video films
E learning programs, Outpatient department, Theaters.

Teaching / Learning Strategies:

Lectures, Small group discussion, Demonstration, Writing Reports, Case Study , e learning ZOOM, GOOGLE MATE etc.

Methods of evaluation and distribution of marks:

*** Quizzes (10 marks)**
*** End of Clinical Course (10 marks)**
**** While midyear exam, Final Clinical Exam & Final written Comprehensive Exam as part of surgical examination**

Recommended textbooks:

1-Operative Surgery Rob & Smith's (Thoracic surgery, Vascular surgery & Cardiac surgery.
2-Mastery of Cardiothoracic Surgery
3-Oxford Handbook of thoracic & Vascular surgery



Pleura is a fibro elastic membrane lines by squamous epithelial cells .
It consists of two portions:-
1-Parietal pleura which lines the thoracic cavity, and it is divided into four parts
a- Costal pleura which lies against the ribs &intercostal muscles
b-Diaphragmatic pleura which covers the upper surface of the diaphragm.
C-Mediastinal pleura which lies against the mediastinum .
D-Cervical pleura which covers the dome of the thoracic space

2-Visceral pleura which intimately invests the lung .



Costo –phrenic angle is the angle between the costal &diaphragmatic pleura.

Cardio –phrenic angle is the angle between the heart &diaphragmatic pleura

Inferior pulmonary ligament is the anterior & posterior reflection of the pleura between the root of the lung & the diaphragmatic surface

The function of the pleura is to maintain the environment of the pleural space in which the lung is function.

Diseases of the pleura
-Spontaneous pneumothorax primary & 2ndary





Is the accumulation of air inside the pleural cavity , occurring without any known etiology .More in males ,more on the right side .It can be bilateral

Clinical Features

Chest pain , pleuritic , sharp in the 1st day then dull pain in 2nd day, cough
dyspnea most common sign & symptom more severe with tension pneumothorax
orthopnea .

Physical examination

may be normal
increased percussion note on the affected side
diminished breath sound on the affected side



Chest X-Ray shows

Pneumothorax

Absence of the lung markings

Shrinkage of the lung towards the hilum & possibly mediastinal shift

Causes

1- Ruptured pulmonary bleb.

2-Ruptured of a cystic defect in the pleura.

3-Teared visceral pleura

4-No cause can be demonstrated in (15-20%).

Complications:-

1-pleural effusion

2-empyema

3-tension pneumothorax which leads to mediastinal shift &circulatory collapse.

4-Respiratory failure in elderly patient with COAD .

Treatment :-

1-Bed rest ,O2 administration &observation in limited pnemothorax.

2-Aspiration

3-Chest tube (thoracostomy tube or ICD intercostal drain in a safety triangle which is bounded by pectoralis muscle anteriorly &lattismus muscle posteriorly and the superior border of the nipple.in the fifth intercostal space just anterior to the mid axillary line to avoid the long thoracic nerve .

4-bronchoscopy is indicated if the lung fail to expand

5-Chemical pleurodesis.by injecting sclerosing agent as Tetracycline, Bleomycin & Talc

6-Surgery pleurectomy by thoracotomy or thoracoscopically if the lung fail to re expand

Up to date 2020

Classification & management of pneumothorax according to Etiology

Pneumothorax type	Specific diagnostic or management strategies to be considered
Primary spontaneous pneumothorax	Likely benign course with conservative management; drainage of pleural gas (typically aspiration), VATS for PAL; lower risk of recurrence.
Secondary spontaneous pneumothorax	PAL is more likely; early intervention with pleurodesis (blood, chemical, surgical) is typically needed; higher risk of recurrence.
COPD	Smoking cessation.
CF	May consider limited pleurodesis strategies if transplantation is planned.
Malignancy	Chemotherapeutic agents or radiation may be appropriate. Pneumothorax may not heal and PAL may be likely such that aggressive surgical strategies may fail.
Infection	Antimicrobials are warranted. Pneumothorax may not heal and PAL may be likely such that aggressive surgical strategies may fail.
Cystic lung disorders	Investigations or therapies targeted at suspected cause may be warranted (eg, lung biopsy, VEGF-D levels, folliculin gene analysis, rapamycin*).
Catamenial (endometriosis)	Hormonal therapy may be warranted.
Architectural abnormalities (eg, Marfan syndrome, Ehlers-Danlos syndrome, Homocystinuria)	May need specific investigations targeted at suspected cause (eg, homocysteine levels).
Iatrogenic	Likely benign course (unless patient is mechanically ventilated). Conservative management with drainage of air is usually sufficient.
Traumatic	May need to co-manage parenchymal trauma and other vascular and orthopedic aspects of chest trauma.
Miscellaneous	
Anorexia	Nutrition needs to be addressed; PAL may be likely.
Exercise	Likely benign course and conservative management with drainage of air may be sufficient.
Illicit drug use	Cessation of drug use.
Immunosuppressant drugs	Cessation of offending agent, if feasible.
Air travel	Avoidance of air travel for short period after definitive management.
Scuba diving	Avoidance of scuba diving until definitive management.

VATS: video-assisted thoracoscopic surgery; PAL: prolonged (persistent) air leak; COPD: chronic obstructive pulmonary disease; CF: cystic fibrosis; VEGF-D: vascular endothelial growth factor-D.

Rapamycin, as an immunosuppressant, is a useful therapy for some patients with lymphangioleiomyomatosis but should not be started until the pneumothorax has healed for about six weeks.

2-Spontaneous haemothorax

Is the presence of blood inside the pleural cavity

Causes:-

1-pulmonary causes ----------TB , AV malformation

2-pleural causes ----------torn vascular adhesion

3-pulmonary malignancy ….primary or metastatic

4-blood dyscrasia ……..hemophilia

5-abdomina; pathology haemo peritoneum

6-thoracic causes ……ruptured great vessels

Clinical features

dyspnea , chest pain ,syncope

signs of hypovolaemic shock

blood inside the pleural cavity may leads to deposition of fibrin on the pleural surface leading to fibrosis (trapped lung syndrome) .

Treatment

1-Resuscitation

2-Tube thoracostomy

3-May needs thoracotomy if excessive bleeding

initial bleeding more than 20ml/kg

Or continuous bleeding more than 3 ml/kg/hour for more than 3 or 4 hours

3- Chylo–thorax

Is the presence of lymph in the pleural space

Causes

A-Congenital atresia of the thoracic duct , birth trauma

B-Traumatic

C-Neoplastic malignancy

D-Infection TB

Diagnosis milky pleural effusion that does not clot and contains fat , fat soluble vitamins & antibodies

Treatment

1-Conservative consists of insertion of tube thoracostomy to drain the effusion , correction of the fluid and electrolytes with nutritional supplement.

2-Surgery consists of ligation of the thoracic duct if the effusion continues for more than two weeks .

4-Pleural effusion

Is the accumulation of fluid in the pleural space from excessive transudation or exudation of the interstitial fluid from the pleural surface. It is signified pleural or systemic disease .

Its effect depends on its size (mild , moderate or massive) & the state of the underlying lung .It is classified as transudate when the protein content is less than 3g% or exudates when protein content is more than 3 gm % .

Clinically patients will present with dyspnea & pleuritic chest pain

Radio logically (concave meniscus sign)

Transudate as in CHF

Exudate as in malignancy

Treatment :-1-aspiration (thoracentesis) 2-tube thoracostomy

3-chemical pleurodesis 4-pleurectomy to remove the pleura to stop the effusion.

5-Empyaema

Is the accumulation of pus in the pleural space , it passes into three stages

1-Acute phase with the clinical manifestation of fever & toxicity .

2-Transitional phase with the increased turbidity of the fluid & decrease the size of the lung .

3-Chronic phase with the pleural thickening ,decrease amount of the fluid & the development of the trapped lung syndrome .

Etiology

1-Secondary to a pneumonic process in the lung.

2-Secondary to a sub phrenic abscess

3-Secondary to a F.B. inhalation .

4-Secondary to esophageal perforation

5-As a post operative complication

Clinical features

Patient may be toxic , febrile , dyspneic with severe cough .CXR shows pleural effusion.

The pus may erode through the bronchus leading to broncho pleural fistula causing severe cough with the production of foul-smelling sputum or the empyaema may presents as a subcutaneous abscess (Empyaema necessitates) which will drain spontaneously .

Diagnosis :-

1-Aspiration of a sample of pus for culture & sensitivity test .

2-Bronchoscopy to exclude TB or malignancy .

Treatment

1-Aspiration of the pus (thoracocentesis) in encysted cases .

2-Tube thoracostomy to drain the pus and to allow the lung to re expand .

3-Surgery (decortication) to remove the parietal and visceral pleura so giving a chance for the lung to re expand .

Surgery in pulmonary tuberculosis





Indications

1-Massive or recurrent haemoptysis 2-Broncho pleural fistula

3-Open cavity , + ve sputum with failure of medical treatment after 3-6/12.

4-Symptomatic patients with destroyed segment or lung & - ve sputum.

5-TB bronchiectasis .

6-When malignancy is suspected .TB & malignancy can co exist or carcinoma can arise in TB scar (Scar carcinoma).

7- Patient with (trapped lung syndrome) after chronic empyaema

Surgery includes <u>segmentectomy , lobectomy ,pnemonectomy & decroticatio</u>ns.

Pleural tumors

Classified as primary & secondary tumors

Primary pleural tumors are **Mesothelioma** which may be of two types

1-Localized benign

2-Diffuse malignant

In both types fibrous , fibro sarcomatous & epithelioid varieties are seen .

Malignant mesothelioma causes chest pain ,bloody pleural effusion with a chest X-ray findings of thickened pleura with nodularity and a limited pleural effusion . There is a possible history of asbestos exposure .

Metastases is uncommon .Death usually occurs within 1-2 years .It has poor response to surgery , radiotherapy & chemotherapy .

Pleural involvement by a metastatic disease is more common than a primary tumor and usually comes from the lung , breast & stomach .

THANK YOU

Chest trauma

Assisted professor

Laith Sharba

MBChB FICMS EACTS FACS

Medical College – JIHMU

Consultant Cardiothoracic and Vascular surgeon

The chest wall, defined here as the bony and muscular structures covering the entire thoracic cavity, protects internal thoracic organs (heart and lungs), mediastinal structures (esophagus and trachea), and major vasculature (aorta and vena cava). Damage to the chest wall may coincide with significant injury to certain of these internal structures and thus, warrants careful evaluation

Chest wall injuries that are associated with increased risk of significant intrathoracic or intraabdominal injury include:

- Scapula fracture
- Flail chest
- Multiple rib fractures (≥3) and displaced rib fractures
- Sternal fracture
- Posterior sternoclavicular dislocation

- **Physical findings of chest wall trauma that are consistent with intrathoracic or intraabdominal injury include:**

- Hypoxia or signs of respiratory difficulty (e.g. , accessory muscle use)
- Paradoxical movement of a chest segment during respiration (suggesting flail chest)
- Hemodynamic instability
- Seat belt sign across the chest or abdominal wall
- Tenderness over multiple rib segments (suggesting multiple rib fractures)
- Palpable depression or deformity of sternum or multiple rib segments
- Abdominal tenderness or guarding

Notes in resustation:

- **Chest compressions should be delivered at a rate of 100 to 120 per minute** because compressions faster than 120 per minute may not allow for cardiac refill and reduce perfusion.
- Chest compressions should be delivered to **adults at a depth between 2 to 2.4 inches (5 to 6 cm)** because compressions at greater depths may result in injury to vital organs without increasing odds of survival.
- Chest compressions should be delivered to **children (less than one-year-old) at a depth of one third the chest, usually about 1.5 to 2 inches (4 to 5 cm).**
- Rescuers must allow for full chest recoil in between compressions to promote cardiac filling.
- Because it is difficult to accurately judge quality of chest compressions, an audiovisual feedback device may be used to optimize delivery of CPR during resuscitation.
- Interruptions of chest compressions, including pre- and post-AED shocks should be as short as possible.
- **Compression to ventilation ratio remains 30:2 for an individual without an advanced airway in place.**

Causes :-

1-Blunt Trauma
2-Pentrating , perforating & blast injuries
3-Iatrogenic injuries

Classification

1-Chest wall injury :

Superficial chest wall injury affecting the skin and wall without affecting the muscles of the chest ribs or the underlying pleura as proved by normal chest x-ray .

Management is to arrest any bleeding and wound debridement with primary closure if possible .

2-Ribs & Sternal injuries :

May lead to

Single rib fracture the important thing is to relief pain by

analgesic , chest wall strapping or intercostal . nerve block

or Multiple ribs fractures , flail chest in that case four or more ribs are fractured anteriorly and posteriorly which lead to paradoxical respiratory movement with the resultant hypoxia and severe dyspnea that may need emergency intubation and assisted ventilation and if the patient needs prolonged intubation for more than one week then tracheostomy is indicated .



End tracheal tube

Flail chest

Costo chondral disruption

hematoma

3-Pleural injuries:

Can lead to

A-Pneumothorax which can be partial or complete
closed or open sucking
or tension pnemothorax .

B-Haemothorax may be mild or severe and may be with pneumothorax





Fractures



Traumatic haemothorax (Localized effusion)

4-Pulmonary injuries:

Trauma can lead to pulmonary contusion with interstitial edema which may lead to consolidation of the lung tissues and can be manage by antibiotics administration and clearing of secretion

5-Tracheo-bronchial injuries :

It may lead to crushing or complete tracheal separation which needs immediate air way management and repair of the trachea.

Bronchial injuries can be so severe that it lead to complete separation of the bronchus which lead to pneumothorax with severe air leak (massive air leak) , hemoptysis &hemothorax .Surgery is mandatory to reanastomosed the bronchus .



6-Great vessels injuries:

Injuries to the thoracic aorta and its branches can occur mainly with deceleration injury . Most patients with ruptured aorta die immediately but in 10% of patients the per aortic tissues and pleura can maintain the intravascular pressure producing false aneurysms ,which can be diagnose by CT angiography and needs immediate surgery .

Traumatic aortic disruption is a time-sensitive injury requiring rapid and accurate diagnosis to prevent death.



Widen mediasinum



Descending aortic injury



Endo vascular stent control

7-Diaphragmatic injuries :

Trauma can lead to rupture of the diaphragm ,most commonly the left side affected ,leading to herniation of the viscera .Stomach is the most frequent organ to herniated followed by the transverse colon , spleen leading to collapse of the lung and mediastinal shift which can be seen on chest X-ray .Barium study is of great help in the diagnosis .Treatment surgery through abdominal , thoracic ,combined approach or by laparoscopy .









8-Esophageal injuries:

Thoracic esophagus rarely injured but mostly the cervical esophagus due to penetrating injury to the cervical region .Pain and dysphagia ,sometime fever .Diagnosis can be established by contrast study which will visualize the site of perforation. Treatment may be conservative by intravenous fluid , heavy antibiotics & NBM for 5-7 days which may be enough to seal the perforation , otherwise surgery is indicated to close the perforation .

9-Cardiac injuries :

May vary from superficial laceration to transmural damage which lead to atrial or ventricular septal defect or coronary artery injury .The patient may present with dyspnea ,hypotension & tachycardia .Diagnosis can be accomplished by Echocardiography .

Management includes resuscitation & immediate thoracotomy or sternotomy to treat the injury .

"The surgeon who should attempt to suture a wound of the heart would lose the respect of his surgical colleagues" - Theodore Bilroth, 1882



Stab to the pericardium - Resuscitative thoracotomy

Ultrasound (FAST exam) — Focused Assessment with Sonography for Trauma (FAST) is an essential part of the primary circulation survey for unstable patients, in whom it often determines management . FAST is used primarily to detect pericardial and intraperitoneal blood, and it is more accurate than any physical examination finding for detecting signs of intra-abdominal injury. In hemodynamically stable patients, FAST can be delayed until the secondary survey and is ideally performed by a second operator while the remainder of the secondary survey is completed

Thoracotomy in chest trauma

90% of chest injuries can be treated by chest tube (tube thoracostomy or intercostal drain (ICD)).

Thoracotomy may be …Immediate (resuscitative thoracotomy)

….Emergency

.. Elective

Emergency Department Thoracotomy

Resuscitative thoracotomy

Penetrating thoracic injury

- Traumatic arrest with previously witnessed cardiac activity (pre-hospital or in-hospital)

- Unresponsive hypotension (BP < 70mmHg)

Blunt thoracic injury

- Unresponsive hypotension (BP < 70mmHg)

- Rapid exsanguinations from chest tube (>1500ml) Drain: ؟؟؟؟

Initial: more than 20ml/kg

1st 3 hours: more than 3ml/kg/hr



Immediate(urgent) thoracotomy

1-Massive haemoptysis

2-Massive haemothorax

. Drain:

Initial: more than 20ml/kg

1st 3 hours: more than 3ml/kg/hr

3-Ruptured bronchus with massive air leak

4-when great vessels injury is suspected

5- In diaphragmatic injury

6-Some cases with esophageal injury .

7-In cardiac injury

Conclusion: Massive air leak and massive haemothorax



Elective thoracotomy

1-Clotted haemothorax

2-Trapped lung syndrome

3-Big foreign body or near vital organs

Complications of chest injuries:

1-ARDS: Syndrome of acute RF with the formation of a non –cardiogenic pulmonary edema leading to reduced lung compliance and hypoxaemia which is refractory to oxygen therapy characterized by bilateral diffuse pulmonary infiltrate on chest X-ray (white lung) .A less severe case (ALI=acute lung injury) which consists of a nonspecific pathological changes in the lung in response to a specific insult .

2-Atelectasis

3-Infection

4-Pulmonary embolism

5-Air embolism

6-Traecheal complications

7-Cardiac arrhythmias



29 10 2007

29 10 2007

29 10 2007

Poly trauma Team Members

Team leader•

- •Anesthetist + assistant
- •General surgeon
- •Orthopedic surgeons
- •A/E physician
- •Two (2) nurses
- •Radiographer
- •Scribe (nurse or doctor)

Our team

Number?

Speciality?

Believe?

Sources?

Thank
You

Lung Diseases

2013

Dr. Laith Sharba

Consultant thoracic surgeon

Assistant Professor

MBChB FICMS(CTS) FACS

وزارة التعليم العالي والبحث العلمي

Ministry of Higher Education & Scientific Research

Pulmonary Echinococcosis (Hydatid Cyst)



Hydatid cyst means cyst full of water .
Hydatid disease of the lung is caused by the small tape worm (Taenia Echinococcus) or (Echinococcus Granulosis) .
Echinococcosis is caused by infection with the tapeworm *Echinococcus*, which belongs to the family Taeniidae. Four species of *Echinococcus* cause infection in humans. *E. granulosus* and *E. multilocularis* are the most common, causing cystic echinococcosis and alveolar echinococcosis, respectively.
The two other species, *E. vogeli* and *E. oligarthus*, cause polycystic echinococcosis but have rarely been associated with human infection.

DPDx

Cystic Echinococcosis

Echinococcus granulosus sensu lato

CDC


1 Adult in small intestine
Definitive Host
Ingestion of eggs (in feces)
Intermediate Host
2 Embryonated egg in feces
2
3 Oncospheres relased
4 Hydatid cysts in various organs
3 Oncosphere hatches; penetrates intestinal wall
4 Hydatid cyst in various organs (commonly liver and lung)
Infective stage
Diagnostic stage





The cyst consists of a germinal layer & cyst fluid containing broad capsule & scoleses . A cellular white hyaline layers are laid down outside the cyst so that the cyst is enclosed by a laminated cyst membrane .As the cyst enlarged , it usually reaches the pleural surface . Compression of the lung tissues produces a thin fibrous layer of atelectatic lung tissue around the cyst (capsule , pericyst or adventia)



Hydatid cyst

Clinical Manifestation

A-Asymptomatic

Any smooth homogenous opacity of uniform density with clear cut border and little or no reaction around it on a chest X-Ray is a hydatid cyst unless proved the other wise .



B. Symptomatic

- **Cough & hemoptysis due to rupture of the cyst , or it can lead to severe dyspnea due to pressure effect, or asphyxia ,or a hypersensitivity reaction ,**
- **If the cyst get infected ,it will lead to formation of lung abscess or bronchiectatic changes .**

Anaphylaxis: rapid overview of emergency management

Diagnosis is made clinically:

The most common signs and symptoms are cutaneous (eg, sudden onset of generalized urticaria, angioedema, flushing, pruritus). However, 10 to 20% of patients have no skin findings.

Danger signs – Rapid progression of symptoms, respiratory distress (eg, stridor, wheezing, dyspnea, increased work of breathing, persistent cough, cyanosis), vomiting, abdominal pain, hypotension, dysrhythmia, chest pain, collapse.

Radiological Findings





1-Smooth homogenous opacity (Intact H.C).

2-Partial rupture (per vesicular pneumocyst).

3-Complete rupture (Water –lilly sign) .

4-Formation of lung abscess(Air –fluid level) .

5-Completely coughed out cyst(empty cavity)

6-Rupture into the pleura (hydropneumothorax)



Serology — Serologic tests are more reliable for diagnosis of E. multilocularis infection than for E. granulosus infection; sensitivity and specificity rates are 95 to 100 percent .

A specific E. multilocularis antigen such as the affinity purified Em2 antigen.

Diagnostic judgment must take into consideration the limitations of serologic testing. The likelihood of a positive serology depends on cyst location and viability. Patients with liver cysts are more likely to be seropositive than patients with lung cysts. Serologic assays are less likely to be positive in the setting of calcified or nonviable cysts. In addition, the sensitivity and specificity of serology is greater for *E. multilocularis* than for *E. granulosus*.

WHO stage	Description	Stage	Size
CE1	Unilocular unechoic cystic lesion with double line sign	Active	<5 cm
			>5 cm
CE2	Multiseptated, "rosette-like" "honeycomb" cyst	Active	Any
CE3a	Cyst with detached membranes (water-lily sign)	Transitional	<5 cm
			>5 cm
CE3b	Cyst with daughter cysts in solid matrix	Transitional	Any
CE4	Cyst with heterogenous hypoechoic/hyperechoic contents; no daughter cysts	Inactive	Any
CE5	Solid plus calcified wall	Inactive	Any

WHO: World Health Organization; CE: cystic echinococcis;

Acute management:

The first and most important treatment in anaphylaxis is epinephrine. There are **NO absolute contraindications to epinephrine** in the setting of anaphylaxis.

A. Airway

Promptly and simultaneously, give:

IM epinephrine (1 mg/mL preparation) – Give epinephrine 0.3 to 0.5 mg IM in the mid-outer thigh. Can repeat every 5 to 15 minutes (or more frequently), as needed. If epinephrine is injected promptly IM, most patients respond to 1, 2, or at most, 3 doses. If symptoms are not responding to epinephrine injections, prepare IV epinephrine for infusion (refer to below).

Place patient in recumbent position, if tolerated, and elevate lower extremities.

Oxygen – Give 8 to 10 L/minute via facemask or up to 100% oxygen, as needed.

Normal saline rapid bolus – Treat hypotension with rapid infusion of 1 to 2 liters IV. Repeat, as needed. Massive fluid shifts with severe loss of intravascular volume can occur. Monitor urine output.

Albuterol (salbutamol) – For bronchospasm resistant to IM epinephrine, give 2.5 to 5 mg in 3 mL saline via nebulizer, or 2 to 3 puffs by metered dose inhaler. Repeat, as needed.

Treatment



B. Medical:

1. **Unfit for surgery**
2. **Metastatic wide-spread**
3. **Stages CE5 (some even CE4)**

Stage CE1 and CE3a cysts <5 cm may be treated with albendazole alone

Stage CE1 and CE3a cysts that are >5 cm may be treated with albendazole in combination with PAIR.

Stage CE2 and CE3b cysts either a modified catheterization technique (eg, non-PAIR percutaneous therapy) or surgery (with adjunctive drug therapy)

Stages CE4 and CE5 are inactive cysts that may be managed with observation



Antiparasitic therapy :

Antiparasitic therapy may be used for **definitive** management in selected cases; it is also a useful **adjunctive** therapy to surgery and percutaneous treatment. Albendazole is the primary antiparasitic agent for treatment of E. granulosus . Albendazole is poorly absorbed and should be ingested with food, ideally with a fatty meal to increase bioavailability (15 mg/kg/day, divided into two doses, to maximum 400 mg orally twice daily with food).

In the absence of albendazole, or for patients with severe adverse reactions to albendazole, mebendazole may be used as an alternative therapy; it is less well absorbed than albendazole .

Albendazole inhibits microtubules assembly, leading to impaired glucose absorption and causing glycogen depletion followed by degeneration of the endoplasmic reticulum and mitochondria of the germinal layer, leading to cell death . Cumulative experience with albendazole suggests that treatment leads to cyst resolution in up to **30 %** of patients
size reduction in another **30 to 50 %**
no change in **20 to 40 %**

Praziquantel has been shown to have protoscolicidal activity, although the efficacy of praziquantel in clinical studies is variable, and, thus far, it does not have a definitive role for primary drug therapy . Praziquantel has been used alone and in combination with albendazole. Limited case series suggest the combination of albendazole and praziquantel may be superior to albendazole alone.

PAIR

Puncture, aspiration, injection, and reaspiration

PAIR is performed under ultrasound or computed tomography (CT) guidance. Following cyst puncture, aspirated fluid should be evaluated histopathologically for presence of protoscolices. Following injection of the protoscolicidal agent, reaspiration should be performed after at least 10 to 15 minutes. The presence of protoscolices in the reaspirated fluid should prompt repeat protoscolicidal injection.

Cysts in extra-abdominal sites such as lung or bone are less amenable to management with PAIR.

Treatment



A. Surgical:

1-Inoculation means to remove it intact.

2-Aspiration &evacuation technique

3-Wedge resection or excision of the cyst with adjacent lung tissue.

4-Segmentectomy ,Lobectomy or Pneumonectomy (rare).




Consultant Laith Sharba
27 6 2007

استشاري جراحة الصدر د.ليث شربه
Innoculation of HC
left lower lobe of the lung
Made with
VideoShow

Ruptured hydatid cyst surgery



Thank you

Lung Diseases

Dr. Laith Sharba
Consultant thoracic surgeon
Assistant Professor
MBChB FICMS(CTS) FACS

Bronchiectasis

Definition

Bronchiectasis is a chronic condition where the bronchi (airways) become abnormally and irreversibly dilated, leading to chronic cough, thick sputum production, and recurrent infections. This dilation results in inefficient mucus clearance, fostering an environment for bacterial growth and chronic inflammation.

Etiology

The condition can arise from multiple causes, including:

- **Infections**: Severe lung infections like pneumonia, tuberculosis, and nontuberculous mycobacterial infections.
- **Immune Deficiencies**: Conditions like common variable immunodeficiency impair the body's ability to fight infections, making patients more prone to developing bronchiectasis.
- **Airway Obstruction**: Blockages, such as foreign bodies or tumors, can lead to bronchiectasis.
- **Allergic Bronchopulmonary Aspergillosis (ABPA)**: A hypersensitivity reaction to *Aspergillus* fungus can cause central bronchiectasis.

Pathophysiology

Bronchiectasis follows a cycle of infection, inflammation, and airway damage:

- **Chronic Inflammation**: Infectious or non-infectious triggers lead to persistent inflammation in the bronchi.
- **Airway Dilation**: Repeated inflammation damages the bronchial walls, causing them to widen and lose their elasticity.
- **Mucus Accumulation**: Damaged airways can't clear mucus effectively, promoting bacterial colonization and recurrent infections, which further worsen the cycle.



Clinical Features

- **Chronic Cough**: The hallmark symptom, often accompanied by large volumes of thick, sometimes purulent, sputum.
- **Recurrent Respiratory Infections**: Patients frequently experience infections that may require antibiotic treatment.
- **Hemoptysis**: In advanced cases, patients may cough up blood.
- **Shortness of Breath and Fatigue**: These symptoms arise due to progressive airway obstruction and lung damage.
- **Wheezing**: Some patients may have wheezing, mimicking asthma or COPD.

Diagnosis

- **High-Resolution CT (HRCT) Scan**: This is the gold standard for diagnosing bronchiectasis. It shows bronchial dilation and thickened airway walls.
- **Sputum Culture**: Identifies pathogens causing infection, particularly chronic ones like *Pseudomonas aeruginosa*.
- **Pulmonary Function Tests**: Evaluate the degree of airflow obstruction and assess lung function.



Extensive Left Lung Bronchiectasis CXR







HRCT shows clustering of markedly dilated airways in both upper lobes (arrow)

Chest CT shows

A. left lower lobe dilated and thickened airways (arrow).

B. Arrowhead shows clusters of airways forming cysts destroying lung



Cylindrical Bronchiectasis in the right middle lobe **(Right Middle Lobe Syndrome)**



Post-radiation traction bronchiectasis (Para mediastinal radiation fibrosis, with loss of volume and mild cylindrical bronchiectasis)



CT scan of Chest with Cystic Fibrosis



Management

1. **Non-Pharmacologic Therapies**:
   - **Airway Clearance Techniques**: Essential for all patients, these techniques (chest physiotherapy, oscillatory positive expiratory pressure devices) help remove mucus and reduce infection risk.
   - **Smoking Cessation and Avoidance of Irritants**: Patients should avoid smoking and other lung irritants to reduce disease progression.
   - **Pulmonary Rehabilitation**: A structured program improves exercise capacity and overall quality of life.


End Expiration
PEEP too low and alveoli collapse/atelectasis decreasing oxygen diffusion (less surface area) and more pressure will be needed to re-expand alveoli leading to atelectotrauma.
Optimal PEEP: alveoli remain open (more alveolar surface area), facilitating oxygen diffusion and less pressure needed to expand the lung.
End Inspiration

2. **Pharmacologic Therapies**:
   - **Antibiotics**: Used to treat active infections and as preventive measures for recurrent exacerbations. For example, macrolides like azithromycin are used to reduce flare-ups, especially for patients without *Pseudomonas* infection.
   - **Mucolytics and Hydration**: Nebulized hypertonic saline helps thin and mobilize mucus for easier expectoration.
   - **Bronchodilators**: These may be prescribed for patients with airway hyperreactivity or wheezing, although they are not universally indicated.

3. Surgical Intervention:

A. Indications for Surgery in Bronchiectasis

The goals of surgery are to remove the affected parts of the lung that are causing symptoms or to prevent complications, such as recurrent infections, hemoptysis, or lung abscesses. Here are the key indications:

1. Localized Bronchiectasis:

2. Recurrent and Severe Infections:

3. Massive Hemoptysis:

- Massive or life-threatening hemoptysis (coughing up large amounts of blood) is a critical indication for surgery. Hemoptysis occurs when bronchial arteries supplying the diseased areas are eroded. If the bleeding cannot be controlled by other means, such as bronchial artery embolization, surgery is indicated to prevent fatal bleeding.

4. **Failure of Medical Management**:
   - This includes cases where quality of life is significantly impacted by persistent symptoms like chronic cough, large amounts of sputum production, and ongoing infection.
5. **Complications of Bronchiectasis**:
   - **Lung abscess**: If the infected area of the lung cannot be drained or treated effectively with antibiotics.
   - **Severe pleurisy or empyema**: In cases of infection spreading to the pleural space.
6. **Non-functioning or Destroyed Lung Segment.**
7. **Prevention of the Spread in Localized Disease**.

B. Types of Surgical Management of Bronchiectasis:

Once the decision for surgery has been made, the specific type of surgery depends on the extent of the disease, the patient's overall health, and the location of the affected lung tissue. Here are the main surgical options:

1. **Lobectomy**: This is the most common surgical procedure performed for bronchiectasis.
2. **Segmentectomy**
3. **Pneumonectomy**: It is reserved for extreme cases where bronchiectasis is widespread and has caused irreparable damage to the lung, particularly when the patient has recurrent infections or massive hemoptysis.

4. **Bronchial Artery Embolization (BAE)**: often used as a temporary or adjunctive procedure in patients with massive hemoptysis. It involves embolizing (blocking) the bronchial arteries that are bleeding, which can stabilize the patient before a more definitive surgery like a lobectomy.

5. **Lung Volume Reduction Surgery (LVRS)**: when the bronchiectasis is associated with other conditions like emphysema, LVRS may be used to remove diseased areas of the lung, allowing the healthier parts to function more effectively.

6. **Minimally Invasive Surgical Techniques (VATS)**

Prevention of Exacerbations:

- **Antibiotic Prophylaxis**: Macrolides like azithromycin are recommended for patients with recurrent exacerbations (two or more per year), as they reduce the frequency of infections and improve symptoms.
- **Inhaled Antibiotics**: For patients with chronic *Pseudomonas* infection, inhaled antibiotics such as tobramycin can help control bacterial load and reduce exacerbations.
- **Vaccination**: Annual flu and pneumococcal vaccines are critical for reducing respiratory infections that can exacerbate bronchiectasis.

Lung Abscess

Definition:

It is a localized area of suppuration and cavitations in the lung. It includes TB , mycotic or parasitic cavitations ,bronchiectasis ,ruptured infected hydatid cyst , even pulmonary infarction with abscess formation & cavitating tumors

Simple lung abscess (pyogenic) can occur as a result of aspiration of a septic debris from the oropharynx into the lung or following dental or tonsillar operation .

Esophageal diseas that lead to regurgitation &subsequent aspiration of esophageal content into the lung is another cause.

Usually, the aspiration is into the RMB leading to severe pnemonitis and liquefaction may occur .As the liquefied necrotic material empties through the bronchus , a necrotic cavity containing pus and air is formed .The organism responsible may be streptococcus , staphylococcus &Escherichia coli .

Clinically cough & foul-smelling sputum , fever , pleuritic chest pain ,night sweat & weight loss .In severe cases dyspnea & cyanosis



Chest X-Ray

Can shows the characteristic air fluid level ,may associated with pleural thickening , pneumothorax

CT chest is helpful in demonstrating the abscess .

Treatment

1- **Primary treatment is** Medical

- **Prolonged antibiotics treatment**
- **Bronchoscopy useful to remove a FB or drainage of abscess**

2-Surgery indicated in

- **Failure of medical treatment**
- **Massive hemoptysis**
- **persistent of a thick wall cavity**
- **when malignancy is suspected**
- **when empyema develops**

Complications :-

- Empyema
- Septicemia
- Metastatic brain abscess
- Bronchogenic spread
- Development of chronicity


Consultant Laith Sharba

Thank You for Your Attention

Endoscopic Thoracic Surgery

Dr. Laith Sharba MD FICMS FACS
Assisted professor
Consultant Cardiothoracic & Vascular Surgeon



Endoscopic Thoracic Surgery

Bronchoscope

Dr. Laith Sharba MD FICMS FACS
Assisted professor
Consultant Cardiothoracic & Vascular Surgeon

Definition

Yesterday: Looking into the living lungs (Chevalier Jackson 1928)

Today : (View the fine details of the end bronchial anatomy ,Diagnosis of the disease & Treating diseases). With the major advance in technology

It is the visualization of the air way using either rigid Bronchoscope (GA) or the flexible (Fiber optic Bronchoscope) (LA) or both simultaneously .Through which we can remove FB , take BAL , brushing lesions & Trans bronchial Biopsy .





HISTORY

Gustav Killian 2 June 1860-24 Feb. 1921 was German Laryngologist **(The father of bronchoscopy) , was appointed at the university of Freiburg in 1892 .**

Gustav Killian in 1897 succeeded in removing aspirated pork bone from the bronchus of a 63 years old farmer under cocaine anesthesia .He used external light source , a head mirror , esophagoscope and forceps to remove the bone .

He became famous & his clinic attracted patients from far and wide for his expertise in removing different kind of (FB) such as bones ,beans ,buttons ,coins & tin whistle .

Gustav Killian Bronchoscope , external light source

Bronchoscopy rapidly developed into a science (with the creation of a better instruments and techniques)

Chevalier Jackson

Founded Philadelphia school of bronchoesophalogology

Jackson‘s monograph first published in 1950



Chevalier Jackson’s Bronchoscope with a small distal bulb & built –in suction tube

Early in the 1960s Shigeto Ikeda devised a means to replace the small electric bulb with glass fibers capable of transmitting brighter light from an outside source. He presented the first flexible bronchoscope at the 1966 International Congress on Diseases of the Chest in Copenhagen.

H.H. Hopkins English physicist developed the rod-lens optical telescopes which could be used with the rigid bronchoscope

At the end of the 1980s, Asahi Pentax replaced the fiberoptic bundle with a charge-coupled sensor at the tip of the scope. This video bronchoscope allowed the bronchoscopist to look at a monitor screen instead of through the eyepiece of the scope

Rigid Bronchoscopes

Hollow metal tubes , of variables sizes down to (3mm - 9.5 mm) and variable length (20-40 cm) .These instruments usually have illumination at their tips as well as side holes near the tip to facilitate ventilation .They are always inserted trans orally. General anesthesia is recommended for its introduction .

Indications for Rigid Bronchoscopy:

1-Hemoptysis

2-Tracheal stenosis.

3-Foreign body removal .

4-All broncho logical interventions such as bougienage , removal of threads & post intervention hemoptysis.

5-Confirmation of cell type in case of previous , non diagnostic fiberoptic bronchoscopy .

6-Laser treatment

7-Removal of tumor

8-Removal of excess fibrin , mucous plug .

9-Bronchography .

10-Autofluorescence and photodynamic diagnosis .

i.e. Diagnostic and Therapeutic



Endoscope for the museum

Bronchoscopy (Adult Set)



Bronchoscopy (Pediatric Set)




19 1:29

PA
ERECT
LEFT





Foreign Body (LMB)

TRACHEOBRONCHIAL FOREIGN BODIES IN PAEDIATRIC AGE GROUP

W. M. Hussein*

Summary

This communication reports the various types of foreign bodies encountered at the Department of Thoracic and Cardiovascular Surgery of the Medical City Teaching Hospital in Baghdad for the past two years. The study analyzes 427 cases. The commonest foreign body extracted was watermelon seed. The clinical manifestations and the radiological appearances of such cases are presented, and also reports what our team has learned from the successful removal of such foreign bodies.

All foreign bodies were successfully removed by bronchoscopy. The procedure of bronchoscopic removal of foreign bodies and the general principles of management are discussed.

Our excellent results are attributed to the early and accurate diagnosis, planned multidisciplinary management, excellent anaesthesia and proper care after the procedure.

Patients and Materials

Definite history of foreign body inhalation could be ob-

* Department of Thoracic and Cardiovascular Surgery, Medical City Hospital, Baghdad, Iraq.

Table 1 : The Different varieties of foreign Bodies in Number :

	Classification	Cases
1.	Watermelon Seed	244
2.	Peanut	55
3.	Sunflower Seed	27
4.	Hazelnut	11
5.	Chicken and fish bones	8
6.	Worry Beads	8
7.	Almond	5

8.	Beans	5
9.	Chick Pea	5
10.	Seeds	5
11.	Apple Pieces	4
12.	Broad Bean	3
13.	Rice	3
14.	Popcorn	3
15.	Clip	2
16.	Pin	2
17.	Pemigranat	2
18.	Tamarind Seed	2
19.	Plastic Pieces	2
20.	Candle Wax	1
21.	Pellets	1
22.	Orange Seed	1
23.	Wood Piece	1
24.	Okra Piece	1
25.	Porcline	1
26.	Toy Piece	1
27.	Egg Shell	1
28.	Screw	1
29.	Metal Piece	1

File Edit View Options

TABLE 51–11. Advantages and Disadvantages of Rigid Bronchoscopy

Advantages	Disadvantages
Durability	General anesthesia
Large instrument channel	Limited distal visualization
Control of airway	Cost
Usefulness in:	
Pediatric bronchoscopy (especially foreign bodies)	
Massive hemoptysis	
Stricture dilation	
Stent placement	
Laser therapy	
Phototherapy	
Cryotherapy	

Flexible Bronchoscopy





It consists of Control section , a flexible insertion tube & a bending tip .The control section contains the eye piece ,control lever and a channel for aspiration or for introduction of solution and instruments.

These flexible bronchoscopes have variable outside diameter (1.8 - 6 mm) with inner channel ranging from (1.6 -2.6 mm) . Highly maneuverable and can reach areas in the endo bronchial tree not accessible to the rigid bronchoscopes .

Can be inserted either trans orally or trans nasally or through the rigid bronchoscope. We have the Infantile , Pediatric , adult types and the Video –Bronchoscopes

Indications for flexible Bronchoscopy

1-For routine examinations .

2-Treatment of acute respiratory problems in the ICU.

3-Suction under visual control .

4-Use of catheter and brushes for cytology.

5-For obtaining secretions for bacteriological tests.

6-Localization of the bleeding site in case of hemoptysis.

7-Theraputic suction & irrigation .

8-Transbronchial lung biopsy .

9-Selective bronchography .

10-Autofluorescence & photodynamic diagnosis .

Major disadvantage of the flexible Bronchoscope is that it is a closed system that does not provide an airway , and the relatively small inner channel is considered to be incapable of allowing adequate suction when confronted with copious secretions or massive hemoptysis .
It is not so much effective in the removal of foreign bodies .

Tracheo-Bronchial Trainer

Suction

Lever

Saline for irrigation

Sample collection (BAL)

Lecture scope

Contra indications

Rigid Bronchoscopy

Is best avoided in the presence of

1. Cervical spine injury : to prevent hyperextension of the neck

2. Aneurysm of the thoracic aorta

Flexible Bronchoscopy

Best avoided in patient with

1. **Massive Hemoptysis**
2. **Air way problems(asthma COPD)**

In cases of doubt as to whether bronchoscopy should be done or not , bronchoscopy should always be done (Jackson's 1915 statement) .

In suspected cases of F.B inhalation it is better to have a negative bronchoscopy rather than to miss a F.B inside with all its pathological consequences .

Preparation

1-Blood tests may be needed before the procedure to ensure that the patient has no problem related to blood clotting as bleeding can occur during or after bronchoscopy specially if biopsies were taken.

2- The patient may be asked to discontinue medications (Aspirin, Plavix ,Warfarin) several days prior to the procedure .

3-The patient should be fasting at least (6-8 hrs.) prior to the procedure .Artificial denture should be removed

Bronchoscopy can be done in a special (Bronchoscopy suite) or operating room or (ICU) .

Anesthesia for Bronchoscopy

Local for the flexible , general for the rigid and combined procedures

Advantages of LA

- Patient is awake
- patient is cooperative???!!!!
- Spontaneous breathing

Disadvantages :

1. Co2 retention may occur
2. Difficult to control bleeding
3. patient may move suddenly (Injuries or damage to scope).

Disadvantage of GA :

1. Patient may not be fit for GA
2. General and specific complication especially cardio- Vascular and ventilation complications .

In both cases the patient should be carefully monitor as regard ECG , BP , and pulse oximetry .

Normal right upper left bronchus.

Normal branching of [redacted] lower left bronchus.

Normal termination of right main bronchus.



Normal left upper bronchus.

Normal Bronchoscopic Findings



Virtual occlusion of left main bronchus by endo-bronchial tumour.



Carcinoma in left main bronchus. Left main occluded by large fleshy tumour, partly obscured by purulent secretions.



Carcinoid adenoma in right intermediate bron-chus. Large red vascular tumour that bled easily and profusely.



Foreign body in right intermediate bronchus.

Abnormal Bronchoscopic Findings

File Edit View Options

TABLE 51–12. Advantages and Disadvantages of Flexible Bronchoscopy

Advantages	Disadvantages
Patient tolerance	Instrument channel size
Topical anesthesia	Sterilization
Field of view	Maintenance
Angle of deflection	
Versatility	
Ambulatory setting	
Ventilator-dependent setting	
Usefulness in:	
Cervical spine disorders	
Subsegmental intervention	
Transbronchial needle aspiration	
Brachytherapy	
Immunotherapy	

Complications of Bronchoscopy

A. Premedication & anesthesia:

Respiratory depression/arrest
Transient hypotension
Seizures
Hyperexcited state
Laryngospasm
Syncope
Cardiac arrest

B. Biopsy:

Hemorrhage
Perforation(Mediastinal emphysema, pneumothorax)
Air embolism

Complications of Bronchoscopy

C. Technical Difficulties:

Respiratory compromise(obstruction, perforation)
Hemorrhage
Laryngospasm
Bronchospasm
Dysrhythmias
Syncope
Pneumonia
Hypoxemia
Pyrexia
Epistaxis

Post bronchoscopy care

1-Close monitoring for 2-4 hours after the procedure

2-Eating and drinking is not allowed until the effect of anesthesia have worn off .

3-Some may advise routine CXR after performing a biopsy to check for signs of pneumothorax .

3-Those patients develop complications may need to stay in the hospital for additional time .

4-The patients may have sore throat , hoarseness ,cough or muscle ache .

Fever up to temperature 38 “ c is common after bronchoscopy but usually for only 24 hours .

Advances in Bronchoscopy

1-Brochoscopic Ultra-sound

2-Bronchoscopic stenting (Air way prosthesis)

3-AF Bronchoscopy .

4-Bronchoscopic Laser therapy .

5-Bronchoscopic Electro Cautery

6-Cryo therapy

7-Brachy therapy

8-Photo therapy

1-Endo bronchial U/S has the ability to visualize via U/S the extra airway structure that cannot be visualized bronchoscopically

Bronchoscopic ultrasound image of the trachea (TR)



2-Bronchoscopic Stenting

Air way prosthesis (Stents) is made of metal (silicone) available in various sizes and shapes .It can be placed in the obstructed air ways to provide relief of symptoms caused by malignant or benign air way disorders

3-Auto florescence Bronchoscopy (AF)

The potentially tumorous areas are darker than the surrounding healthy tissues i.e. AF creates the color contrast required to demarcate potentially tumorous areas (Useful for early detection of Ca .Bronchus)





Mucosal metastasis of a carcinoma, not visible under white light but clearly visible in autofluorescent mode





4-Laser Therapy

Using Nd-YAG laser considered in patient with surgically unresectable lesion .The treatment is aimed at relieving obstructive symptoms rather than curing the neoplasm .It is used to treat both benign and malignant air way tumors that obstruct the major airway also help in preparing the air way for insertion of air way stent, but it can lead to severe hemorrhage , pneumothorax and pneumomediastinum





5-Bronchoscopic Electro cautery:

It employs alternating electrical current to produce coagulation & vaporization of the endobronchial lesion .(Argon plasma coagulator also used). Probes are available to be used with both rigid and flexible bronchoscopes .It leads to relief of symptoms caused by air way obstruction .It is less expensive than laser .
It can lead to:

- End bronchial fire
- Hemorrhage
- Electric shock to the operator or the patient.

6-Bronchoscopic Cryotherapy:

Bronchoscopic cryotherapy consists of cold-induced death of malignant cells by repeated cycles of cold application followed by thawing . Nitrous oxide or liquid nitrogen is most used to produce temperature of -80 C .

Indications are the same, but it is less expensive than laser with fewer complications however it needs longer duration of therapy

7-Brachy therapy :

Brachytherapy is used to deliver radiation therapy from the airway lumen to treat malignant tumors that within obstruct the airways. The role of bronchoscopy is to place the brachytherapy catheter bearing the radio active source into the obstructed air way under direct vision .

8-Photo therapy:

Within 24 to 48 h after IV administration of hematoporphyrin derivative (Florescent substance) , light of an appropriate wavelength is shined on the tumor for several minutes during bronchoscopy. Response lasting for > 12 months has been observed in 50% of patients. May lead to expectoration of thick necrotic material which may needs therapeutic bronchoscopy

Bronchoscopy need cooperation and mutual understanding Between

1-A well trained endoscopist

2-a qualified staff

3-Expert and well-trained anesthetist

Rigid Bronchoscope for FB inhalation



Rigid Bronchoscope for emergency



THANK YOU FOR YOUR ATTENTION

Endoscopic Thoracic Surgery

Dr. Laith Sharba MD FICMS FACS
Assisted professor
Consultant Cardiothoracic & Vascular Surgeon



Endoscopic Thoracic Surgery

slido

Join at slido.com
#3291129

ⓘ Start presenting to display the joining instructions on this slide.

Introduction

Video-assisted thoracoscopic surgery **(VATS)** is a set of minimally invasive thoracic surgical **(MITS)** procedures used to diagnose or treat conditions of the chest (pulmonary, mediastinal, chest wall). Most major procedures traditionally performed with open thoracotomy can be performed using smaller incisions with video assistance. A related technology, robotic-assisted thoracic surgery **(RATS),** uses computers to aid surgeon instrument control.

Thoracoscopic surgery In the same way laparoscopic techniques reduce the need for large abdominal incisions, minimally invasive thoracic surgery (MITS) eliminates the need for thoracotomy that requires spreading of the ribs or sternotomy incisions.MITS uses a thoracoscope attached to a video camera to see into the chest.





Terminology

The use of the term "video-assisted thoracoscopic/thoracic surgery" (i.e., "VATS") is applied to multiple similar, but not identical, methods. There is variability in the placement, size, and number of the incisions (also called ports) and the extent of usage of the thoracoscope that is used to accomplish the surgery.

* VATS uses an access (or utility) incision that ranges from 2.5 to 8 cm in length and allows manipulation of multiple traditional or thoracoscopic instruments through the same incision at the same time. VATS can be performed with one (uniportal) or up to four chest incisions



MITS: Two instruments in one port Video



**"Totally thoracoscopic" or "completely thoracoscopic" or "totally portal" procedures use ports that are only large enough to admit a trocar without the use of an access or utility incision. This facilitates insufflation with carbon dioxide to maintain exposure but may require enlargement of one incision to extract specimens.

slido

Question: In the context of totally thoracoscopic or completely thoracoscopic procedures, which of the following statements is accurate?

ⓘ Start presenting to display the poll results on this slide.

***The other method of minimally invasive surgery in the chest is robotic-assisted thoracoscopic surgery (RATS). RATS uses a unit that has multiple movable arms controlled by the surgeon at a separate ergonomic console as currently there is the only intracorporeal robotic system for use in humans approved for use in the United States. Robotic systems are more properly referred to as Computer-Aided Surgical Systems (CASSs) because there is essentially no robot movement without human guidance.


EndoWrist Instruments are designed with seven degrees of freedom and mimic the dexterity of the human hand and wrist
Patient cart
Anesthesiologist
Vision cart
Assistant
Nurse
Surgeon uses open-surgery hand movements that are precisely replicated in the operative field by the EndoWrist Instruments
Surgeon at console

slido

Question: Which of the following best describes the relationship and differences between Video-assisted thoracoscopic surgery (VATS) and robotic-assisted thoracic surgery (RATS)?

ⓘ Start presenting to display the poll results on this slide.

Comparison between VATS & RATS

Feature	VATS	RATS
Accessibility of equipment, competitive vendors, flexibility of systems	+++	+
Expense	++	+++
Ergonomics	+/–	++
Use of carbon dioxide	+/–	++
Ability to reduce the number of port sites (ie, multiple instruments at a single site)	+++	–
Movement scaling and tremor reduction	–	+++
Ease of conversion to open surgery	++	+
Availability of 3D optics	+	+++
Capability in confined space	+	+++
Replication of open instrument function	+/–	++
Skilled assistant needed	++	+

slido

What is the more proper term for robotic systems used in surgeries like RATS?

ⓘ Start presenting to display the poll results on this slide.

slido

What is a common requirement for both VATS and RATS procedures when it comes to extracting specimens?

ⓘ Start presenting to display the poll results on this slide.

slido

What is a significant difference in the technology used between VATS and RATS?

ⓘ Start presenting to display the poll results on this slide.

KEYS OF SUCCESS
Good Camera Man
Hand Eye Coordination skill
SHOT ON MI NOTE 10
108MP PENTA CAMERA

2K
With
OLD Monitor

2K with upgrade monitor
SHOT ON MI NOTE 10
108MP PENTA CAMERA

SHOT ON MI NOTE 10
108MP PENTA CAMERA

GOOD VISION
4K
SHOT ON MI NOTE 10
108MP PENTA CAMERA

3D for more sophisticated operation

MITS provides safe, effective, and successful surgery when patients are selected appropriately. The indications have expanded as technology has improved. Continued outcome assessments are needed to ensure that MITS provides equivalent or improved outcomes compared with traditional open surgical methods.

In the same manner in which laparoscopic techniques reduce the need for large abdominal incisions, minimally invasive thoracic surgery (MITS) eliminates the need for thoracotomy that requires spreading of the ribs or sternotomy incisions. MITS uses a thoracoscope attached to a video camera to see into the chest.

Thoracoscopic tool


A
B

A

B

C


A
B

Indications

Minimally invasive approaches can be used for the diagnosis or treatment of benign or malignant chest diseases. Many procedures historically performed as an open thoracotomy are now performed as video-assisted thoracoscopic surgery (VATS). As surgeons become more proficient and technological advances in optics and instrumentation occur, the number and complexity of diseases/problems that can be safely and reliably diagnosed or treated by thoracoscopy are growing.

A. Diagnostic

Thoracoscopy was initially used mainly as a diagnostic tool for pleural diseases, particularly for tuberculosis For diagnostic purposes, pulmonologists, general surgeons, trauma surgeons, and thoracic surgeons may use thoracoscopy to visually inspect the structures of the chest or to obtain fluid or tissue for histologic examination or cultures

1) **Pleural biopsy**
2) A variety of other tissues can be accessed for **inspection or biopsy** (e.g., mediastinal nodes, diaphragm, lung parenchyma, pericardium, and esophagus)
3) Also, if malignant pleural disease is discovered, thoracoscopy can **guide** pleurodesis or a placement of a drainage catheter.



Thoracoscopic pleural metastasis

B. Therapeutic

Therapeutic minimally invasive thoracic surgery (MITS) slowly gained momentum in the 1990s after the success of laparoscopic operations such as cholecystectomy. The most common uses for therapeutic MITS include the following:

A) Pulmonary resection

B) Pleural disease/chest cavity : Pleural drainages (pneumothorax, haemothorax, empyema, malignant pleural effusion) with mechanical or chemical pleurodesis are approached commonly using thoracoscopy.

C) Diaphragm surgery : involves plicating or resection of the diaphragm. Diaphragmatic injury can also be evaluated and repaired thoracoscopically

D) Chest wall surgery : chest wall deformities (e.g., pectus), rib biopsy, and chest wall tumor resection

E. Esophageal surgery : MITS for benign diseases of the esophagus and mediastinum allows successful treatment using smaller incisions. Most esophageal pathologies that are amenable to a thoracoscopy (e.g., excision of esophageal diverticulum are approached from the right chest cavity; however, the more distal thoracic esophagus (e.g., myotomy for achalasia and the gastro esophageal junction),distal esophageal carcinoma, leiomyoma can be approached either from the left chest cavity or laparoscopically .



Thoracoscopic diverticulum



Myotomy for achalasia



Thoracoscopic Lower esophageal Tumor



Thoracoscopic Lower esophageal CA

Thoracoscopic Leiomyoma

G) Heart – Cardiac surgeries using thoracoscopy or robotics include coronary artery bypass grafting, atrial septal defect repair, resection of intracardiac tumors, mitral valve repair/replacement, ablation of atrial fibrillation, placement of epicardial pacemaker leads, and creation of a pericardial window

H) Spine – Spine procedures that can be accomplished using a minimally invasive approach include thoracoscopic laminectomy, disc decompression/debridement, and abscess drainage, typically by orthopedic spine surgeons or neurosurgeons.

Contraindication

To adequately see and have space to manipulate instruments within the pleural cavity, it is necessary to collapse the lung on the operative side by selectively ventilating the contralateral lung. The **only absolute contraindications to VATS and RATS are the inability to achieve the working space needed or if a patient cannot tolerate one-lung ventilation** (e.g., airway mass, prior pulmonary resection, severe pulmonary disease).

slido

Which of the following is a disadvantage of minimally invasive procedures (VATS and RATS) compared to open surgery?

ⓘ Start presenting to display the poll results on this slide.

slido

How do the overall costs of minimally invasive procedures compare to open surgery, considering the length of stay and recovery?

ⓘ Start presenting to display the poll results on this slide.

The advantage over open:

*Quality of life assessments
*Morbidity rates
*Recovery timelines

Disadvantage:

perioperative costs for minimally invasive procedures (both VATS and RATS) are higher because of the equipment. However, overall costs may be lower due to shorter length of stay and faster patient recovery

slido

Audience Q&A Session

ⓘ Start presenting to display the audience questions on this slide.

THANK YOU FOR YOUR ATTENTION

DR.HUSSAIN SAHIB
Consultant neurosurgeon
Neurosurgery

Head Injury

External trauma applied to the head including, scalp, skull and it's containing brain tissues, which result of:

- 1-**closed** trauma as in road traffic accident falls or assaults.
- 2-**open** trauma as in gunshots, stab wounds.

Clinical classification of head injury

Primary head injury can be classified into four main categories include:

A-scalp injuries like.

Brusies, wounds, laceration, avulsion and hematoma

B-skull fractures:.

Linear, diastatic, depressed, basilar.

Skull fractures result from large energy force applied to the head during injury; the extent and type of skull fracture depend on the severity, direction and geometry of impact force.

1-linear fracture:

Is a single fracture line in the skull, which passes through the entire thickness of the skull.

Linear fracture indicates that patient received an impact of considerable energy, but the patient eventual neurological disability depends on brain injury not the linear fracture.

It increases the risk of intracranial hemorrhage, for this reason detection of skull fracture in plain x-ray indicates the need for ct-scan of head.

Linear skull fracture require no stabilization if the scalp is intact. An open linear fracture should be inspected, removal of foreign body and debridement of damaged soft tissues.

2-*diastatic fracture*:

Is a separation of suture line, its common in infants under three years old and rare in older age group except as apart of more extension of skull fracture.

3-*depressed fracture*

Result when outer table of fractured edge lies below the outer table of surrounding intact skull at the site of impact, it is either full thickness when the outer table of the depressed segment is below the inner table of the non-depressed segment or partial if above it.
Depressed fracture occur due to more localized force of impact applied to the skull as by hammer, sport injuries like hockey stick, golf ball or vehicle accidents .its either closed (intact scalp) or opened (compound) when connect to the exterior through scalp injury.
Depressed fracture may associate with scalp laceration, dural laceration, brain contusion and intracranial hemorrhage.
Diagnosis is made on the basis of skull x-ray and ct-scan of head with bone windows.
Treatment of individual depressed fracture depends on the presence or absence of:
-----cosmetic deformity.
-----scalp laceration.
-----dural laceration.
-----contusion or laceration of underlying brain.
-----extension of fracture to the air sinuses.
-----coexistence of associated intracranial lesion like EDH, SDH, ICH, cerebral contusion, midline shift and ventricular compression.
Treatment usually in form of elevation of depressed segment in case of closed fracture or removal of depressed segment in case of compound full thickness depressed fracture followed by subsequent cranioplasty using titanium mesh ,acrylic(artificial bone) or bone graft 4_6 month later if the pt is above 4yrs age .

4-basilar fracture:

Which involve anterior, middle, or posterior cranial fossae , result due to either direct impact or propagation of stress wave through the skull from a remote impact.

Clinical sign of basilar fracture include cerebrospinal (CSF) leakage from ear (otorrhea),or nose (rhinorrhea),blood behind the ear drum (hemotympanum) ,bruising behind ear (postauriculr ecchymosis or Battles sign)bruising around eye (periorbital ecchymosis or Raccoon eye) .cranial nerves injury specially facial (7th) or vestibulocochlear (8th) may occur.

The presence of these clinical sign should increase the index of suspicion and help in the identification of basal skull fracture.

Most of basilar fracture require no treatment, persistence CSF leak may require operative repair.

C-focal brain injuries:

Defined as macroscopically visible damage to the brain parenchyma that is generally limited to a well defined area.

Focal brain injury includes:

1-contusion:



Bruises of neural parenchyma, its either coup contusion which occur at site of impact or countercoup when occur remote from site of impact force.

Commonly occur at frontal and temporal area of brain but can occur anywhere.

2-intracranial hematoma:



--EDH --ICH --IVH

--SDH . --SAH

--Extradural hematoma (EDH) :

Collection of blood between the inner table of the skull and dura.it has 3 sources

a) arterial source the commonest and commonly from middle meningeal artery (MMA).

b) venous source i.e venous sinuses .

c) osseous from the skull bone.

EDH associated with skull fracture in 90% of cases.

Clinical manifestations:

Disturb conscious level, Lucida interval (transient loss of consciousness followed by return of conscious level until the growing EDH result in unconsciousness again) ,hemiparesis , pupillary dilatation.
Can be diagnosed by brain ct-scan through which hematoma will appear as a hyperdense, biconvex (lens shape) area between the skull and dura.
Treatment should be directed first towards reducing sign and symptoms of raised intracranial pressure (ICP) by **medical** treatment like head elevation, hyperventilation, hypothermia, hypertonic saline infusion (mannitol) ,furosomide , steroid , and barbiturate .



Surgery in form of craniotomy and evacuation of hematoma.

--**subdural hematoma :**



SDH is a collection of blood in the subdural space between the dura and brain tissue, result from tearing of the cortical vessels into subdural space .

subdural hematoma whether acute , sub acute or chronic hematoma depend on the age of trauma ,first week , second week or more than two Weeks respectively.
SDH appear as crescentic shape in brain ct-scan may or may not associate with skull fracture.
Treatment should be by evacuation of hematoma by craniotomy and coagulation of cortical bleeder vessels (mainly done in acute SDH) or through burr hole evacuation , through which one or more burr hole opening of skull ,then opening the dura and evacuation of hematoma (mainly done in chronic SDH) .

--**intracerebral hematoma (ICH):**

ICH is a collection of blood inside brain parenchyma, the mechanism is usually acceleration –deceleration with shearing of perforating vessels, cortical or subcortical . it is usually round shape ,most commonly in the frontal and temporal region .

--**subarachnoid hemorrhage** (SAH) :

SAH mean hemorrhage in the subarachnoid space due to the rupture of the vessels in that space.

--**intraventricular hemorrhage (IVH) :**

IVH hemorrhage within the ventricular system it is the worst type.



D-diffuse brain injury :

- Concussion.
- Brain edema.
- Diffuse axonal injury.

--**concussion** **:** a transient reversible neurological dysfunction due to trauma .these injuries commonly occur after head injury and presented in many forms : . transient loss of consciousness
. Retrograde amnesia.
. Headache.
.Vomiting.

--**brain edema (brain swelling):**

Brain edema refer to an increase in brain volume caused by increased intracranial blood volume ,most fluid in edema accumulate in white matter in which can expand and thus compressing further brain tissues .

--**diffuse axonal injury (DAI):**

DAI refer to a prolonged post traumatic loss of consciousness rendering the patient deeply comatose with a features of decortication or decerebration. Usually associated with brain stem injury.

So clinical classification of head injury:

Scalp injury	skull fracture	Focal brain injury	diffuse brain injury
Bruises	linear	contusion	concussion
Wounds	diastatic	EDH	brain edema
Laceration	depressed	SDH	DAI
Avulsion	basilar	ICH	
Hematoma		SAH	
		IVH	

Another important clinical grading of H.I for you as a resident doctors In ER depends on GCS as:-

Minimal H.I : GCS 15/15 without period of loss of consciousness.
Mild H.I : GCS 13-15.
Moderate H.I : GCS 9-12
Severe H.I : GCS 8 and below

Best eye response (E)	Best verbal response (V)	Best motor response (M)
4 Eyes opening spontaneously	5 Oriented	6 Obeys commands
3 Eye opening to speech	4 Confused	5 Localises to pain
2 Eye opening in response to pain	3 Inappropriate words	4 Withdraws from pain
1 No eye opening	2 Incomprehensible sounds	3 Flexion in response to pain
	1 None	2 Extension to pain
		1 No motor response



DR.HUSSAINSAHIB
Consultant neurosurgeon
Neurosurgery

Neurological assessment in head trauma

Neurological assessment include several measure used to define the severity of head injury and important lines taken in management including diagnosis and treatment.
Neurological assessment include following:

1-vital signs monitoring.

Blood pressure, pulse rate, respiratory rate and temperature .its important to assess vital sign in head injured patient as disturbed these sign can give clue to the associated intracranial hypertension in head trauma .
For example Cushing reflex (increased blood pressure with associated bradycardia) can occur with severe head injury.

2-state of consciousness:

Consciousness is a state of arousal or awareness level or ability to react with the external environment .it can be affected by any brain insult ,so Glasgow comma scale GCS is indicated in head trauma to assess the degree of head injury and so the conscious level disability.
GCS consist of three main parameters: eye opening, best verbal response and best motor response with a total of 15 points.

Eye opening	4 \ spontaneously open eye 3\ open to speech 2 \ open to painful stimuli 1 \ none
Best verbal response	5 \ oriented 4 \ confused 3 \ inappropriate words 2 \ incomprehensible sounds 1\ none
Best motor response	6 \ obey commands 5 \ localizing to pain



4 \ flexing to pain (flexion withdrawal)
3 \ abnormal flexion
2\ extension
1 \none

3-pupils examination:

Including pupillary size and reaction to light (light reflex).
The aspect optic oculomotor system can be tested by pupillary light reflex , which require the afferent link of optic nerves and tracts be intact as well as the parasympathetic oculomotor outflow for the efferent link.

4-cranial nerves examination:

The twelve cranial nerves evaluation is mandatory as it can be injured by trauma.
Abducent nerve (6th cranial nerve)commonly affected due to long anatomical course in the skull so it may be injured at any area been traumatized by pathology ,olfactory nerve (1st cranial nerve) ,facial nerve (7th cranial nerve) vestibulocichlear nerve (8th cranial nerve) can injured by skull base fracture .

5-motor response:

Motor system examination should be carried out in trauma as patient may have focal weakness or paralysis or hemiplegia.
Motor examination includes examination of bulk, power, tone, reflexes, coordination and gait.In reflexes babinskis reflex is a very imp.one.

6-sensory response:

Sensory system review, including superficial (pain, temperature, tactile sensation) and deep sensation (position and vibration sense).

7-respiration:

Cortical dysfunction due to trauma may result in respiratory irregularities in form of **cheyne-stokes** breathing ,which demonstrate intermixed shallow and deep respiration in cyclical fashion ,which result in overall hypoventilation leading to hypercarbia and hypoxia .
Hyperventilation may occur at pons injury, ataxic or irregular respiration may occur in injury at level of medulla.

8-herniation syndrome (coning):

Two types of coning, transtentorial and central.

transtentorial coning can be in downward direction due to the temporal lobe uncus pressing on brainstem resulting ipsilateral third nerve palsy ,Cushing reflex and decerebrate rigidity ,it can also occur in upward direction in posterior fosse mass or bleed pushing cerebellar tissue up through tentorial incisura.

Central coning occurs when there is herniation of cerebellum and medulla down through foramen magnum resulting in cheyne stokes breathing, apnea and neck stiffness.

8-brain stem death:

Intracranial swelling and brain damage may lead to cessation of all brain function.

9-polysystmatic examination:

Multi systemic review and examination are mandatory as a part of neurological assessment as associated polysystemic injury may be expected like cardiovascular, respiratory, abdominal, musculoskeletal and genitourinary, that's why team work should involve in assessment and management of head injured patient.

Management:

Include radiological evaluation, neurological assessment and treatment of resulting lesion.

-radiology:

- Skull x-ray
- Brain ct-scan
- Brain MRI
- EEG electroencephalogram
- Angiography

-treatment:

Treatment should be carried out at site of accident, during transport, at emergency room, at intensive care unit ICU, at theatre and on ward.

Types of treatment:

1-ABC:

Airway, breathing, ciculation.

2- neurological assessment as we discussed before.

3- ICP intracranial pressure monitoring:

There is evidence that the monitoring of ICP and the management of elevated ICP in comatose brain injured patient may significantly improve the outcome of sever head injuries, as raised ICP will reduce cerebral perfusion.
4- nutritional support, fluid, electrolyte, and metabolic derangement support.
5- medical and surgical treatment of sign and symptoms of increased ICP ,which include the following:

Medical:

a) –**Head elevation**: to 30 degrees can significantly reduce ICP through gravitational drainage of blood and cerebrospinal fluid CSF so reducing the intracranial volume which in turn reduces ICP all of which can improve the outcome.

b) **Hyperventilation**: is one of the most effective means of controlling increased ICP .its achieve it effect by reducing cerebral blood volume and cerebral blood flow through hypocapnea (decrease PCO2) from resulting vasoconstriction since hypercapnia produce cerebral arterial dilatation. Hyperventilation can give through high flow oxygenation by mask, ambubag or through endotracheal intubations and assisted ventilation.

c) **-Hypothermia**: cerebral blood flow and oxygen metabolism have been shown to decline with hypothermia, as increase body temperature can increase ICP, increased cerebral blood volume and cerebral edema.

d) **-Hypertonic solution (**Mannitol): using hyperosmotic or hypertonic drugs to drive the water from brain to the plasma in addition that it results in CSF formation inhibition.mannitol given in a dose 0.5-1g per kg over 30 minutes.

e) **Furosemide**: can reduce ICP through decreasing brain edema and CSF production.

f) **Steroids:** dexamethasone can have beneficial effect in reducing CSF production and so brain edema although there is controversy about its effect in head injury.

g) **Barbiturate:** it act through reducing brain metabolism and cerebral blood flow, with care that high dose barbiturate can have a deleterious effect through respiratory suppression.

Surgical: in form of:

a) **Evacuation** of hematomas like EDH, SDH, ICH by burrhole or craniotomy.
b) **Ventriculostomy** to drain CSF or hemorrhage.
c) **Craniectomy** for associated depressed fracture or extensive brain contusion.

6-Postoperative care, continous monitoring and physiotherapy.

Dr.hussainsahib

Neurosurgery

Hydrocephalus

Hydrocephalus is defined as an imbalance of cerebrospinal fluid CSF formation and absorption resulting in accumulation of fluid within cerebral ventricles.



Physiology of CSF compartment:

Csf is clear colourless ultrafilterate of plasma, bathes the neural tissue, it serves to protect the brain, and also it's in direct communication with extra cellular space of brain and it helps to deliver, circulate and clear substrates of brain metabolism.

CSF is made almost entirely within the ventricles by the choroids plexus, it circulate through ventricles from lateral ventricle into third ventricle through the interventricular foramen of monro then through aqueduct of sylvius into fourth ventricle.CSF exit the ventricular system through the foramina of luschka and megendie to circulate in subarachnoid spaces over the surface of the brain and spinal cord ,then it is absorbed into venous system by arachnoids granulations in the sup.sagittal sinus.

Classification of hydrocephalus:

For clinical purposes it useful to divide defects of absorption into two types:
1-lesions that obstruct the ventricular system leading to obstructive (non-communicating) hydrocephalus.
2-lesions that obstruct the subarachnoid space leading to non-obstructive (communicating) hydrocephalus.

Causes :

Obstructive hydrocephalus:

- Aqueduct stenosis.
- Tumors.
- Vascular malformation.
- Foreign body.

Non obstructive hydrocephalus:

- Infection as meningitis. meningitis → inflamed arachnoid villi → scarring and obliteration of arachnoid villi
- Hemorrhage. hemorrhage → inflammatory response → fibrosis → obliteration of arachnoid villi
- Tumors. Choroid plexus papilloma

- Arnold chiari malformation.



1 2 3

Clinical features of infantile hydrocephalus:

The diagnosis of infantile hydrocephalus is sometimes obvious at birth.
Clinical feature include:

1. Increased head circumference and head appear large, and body appears small in comparison.
2. Failure to thrive.
3. Delayed mile stones.
4. Craniofacial disproportion with low set ears and eyes.
5. Scalp appears thin and glistening with prominent and distended scalp veins.
6. Anterior fontanel typically enlarged, tense, and bulging.
7. Suture diastases.
8. Abducent nerve palsy (6th cranial nerve) either unilateral or bilateral.
9. sun-setting sign due to upward gaze palsy from 3rd ventricular pressure on the mid brain tectum.
10. Positive translumination sign of head.
11. Cracked pot sound on skull percussion.

MacEwen sign (cracked pot sign): percussion of the skull near the junction of the frontal, temporal, and parietal bones evokes an unusually resonant sound

Adult hydrocephalus:

Adult patient with hydrocephalus may present with either:

1. Acute onset and rapid deterioration.
2. Gradual onset and slowly progressive deterioration.

Acute onset adult hydrocephalus .this type of presentation occurs particularly with tumors causing obstructive hydrocephalus, although it may occur with any of the causes of hydrocephalus.
The major presenting features are due to signs and symptoms of raised ICP

- Headache
- Vomiting
- Papilloedema
- Deterioration of the consciousness level.

Gradual onset adult hydrocephalus. this type of onset occurs less frequently than the previous type, the symptoms of raised ICP are only very gradually progressive and late diagnosis is common. early feature in adolescent involve deteriorating school performance as a result of headaches, failing mental function, memory loss, and behavioral disturbance. Endocrine abnormalities such as infantilism or precocious

puberty due to hypothalamic and pituitary disturbances from chronic hydrocephalus. If the condition is unrecognized progressive visual failure will occur secondary to papilloedema and optic atrophy.

In elderly patient a chronic form of hydrocephalus is called normal pressure hydrocephalus (groups of patients with symptoms of dementia, ataxia and incontienence, where radiological studies showed hydrocephalus but normal CSF pressure.

Radiological investigation

The most important investigation is either **CT** scan or **MRI** of the brain which show which ventricles are dilated. If the lateral ventricle and 3rd ventricles are dilated, and the 4th ventricle is small, it's likely that the obstruction is at the level of aqueduct of sylvius. An enhanced CT scan or MRI will help to determine the cause as obstructing tumor. In a communicating hydrocephalus all the ventricles are dilated.

Ultrasonography through the open anterior fontanel is useful in assessing ventricular size in infants and may indicate the need for repeated CT scans.

Treatment:

In general the treatment of hydrocephalus is a CSF shunt. If there has been rapid neurological deterioration this will need to be performed as an emergency.

If the hydrocephalus is due to an obstructing tumor that is surgically accessible, resection of the mass may lead to resolution of the hydrocephalus and shunt might not be necessary.

CSF shunt surgery: they are of 4 main types including:

- VP shunt (ventriculo-peritoneal).
- VAshunt(ventriculo-atrial)
- LP shunt (lumbo-peritoneal).
- Ventriculo-pleural shunt.
-

Operative procedure:

The usual method of CSF diversion is a ventriculoperitoneal shunt, in which a catheter is placed into the lateral ventricle and is connected to a subcutaneous valve which is attached to a catheter threaded subcutaneously down to the abdomen and inserted into the peritoneal cavity. Alternative drainage sites such as atrium, pleural cavity and lumbar area can be used Instead of the ventricle.



Complications of shunt surgery:

- ***Ventriculo peritoneal shunt VP***

1. Infection: shunt infection is a disastrous complication ,particularly in patient with shunt dependency ,so it avoided by meticulous surgical technique and using intraoperative prophylactic antibiotics which can reduce such unwanted complication.
An infected shunt almost needs to be removed and replaced by a new shunt, preferably in a different position and under appropriate antibiotic cover.
2. Obstruction of catheter by infection or kinking.
3. Malpostion of catheter, when not introduce into peritoneal cavity as into preperitoneal fat.
4. Peritoineal infection (peritonitis)
5. Ascitis.
6. Shunt disconnection or transmigration.
7. Itracranial hemorrhage may occur following a VP insertion in form of intracerebral or sub dural hemorrhage.
8. Bowel injury or perforation.



- ***Ventriculo atrial VA shunt:***

VA shunts when shunting the CSF from ventricle into the right atrium through great vein in the neck as internal jugular vein. A shunt used when there has been marked intraperitoneal sepsis or multiple abdominal operations. It possible complications include pulmonary infarction, cardiac injury, endocarditis, thrombosis and embolism.

- ***Lumboperitoneal LP shunt:*** communicating hydrocephalus

LP shunt involves diversion of CSF from spinal subarachnoid space into peritoneal cavity, it considered only in patient with communicating hydrocephalus. possible complications :
1. Subdural bleeding.
2. Sciatic pain from nerve root irritation.
3. Infection.

- ***Ventriculopleural shunt:***

When CSF diversion from the ventricle into pleural cavity, with possible complication like, pnemothorax, pleural effusion and pleural empymea.

Dr.hussainsahib
NEUROSURGERY

Spinal dysraphism (neural tube defects) NTD

Definition: any malformation of developing CNS whether related to the back or head called NTD, it refers to a group of congenital anomalies of the spine or the head in which midline structures fail to fuse.
Clinically it divided into that whether or not they are covered with normal skin into:

1-**Closed NTD**(spina bifida aperta): covered by skin, like:

I. ***Encephalocele:*** congenital defect of cranium in which portion of CNS herniated through defect.



I. ***Lipomeningomyelocele:*** cystic swelling which consist of meninges and contain CSF , neural tissues and fat tissues.

2-**open NTD** (spina bifida cystica) not covered by skin, like:

I. ***Anencephaly*:** an open lesion on the head with an exposed degenerated brain tissue.

II. ***Meningocele:*** cystic swelling contains meninges and CSF with no neural tissues.

II. ***Meningomyelocele:*** cystic swelling contains meninges, CSF, and neural tissues. It is the most common and important form of spinal dysraphism presenting during the neonatal period. It's characterized by the neural element protruding through a vertebral defect into a meningeal lined sac; it may result in irreversible neurological deficit due to the associated spinal cord abnormality.

This disorder occurs in approximately 2\1000 live birth. There is greater frequency among white than black with slight female predominance. There is a familial incidence, if one member of a family is affected the risk of the disorder occurring in subsequent offspring is about 5%.

Prenatal diagnosis of meningomyelocele and other open NTD disorder is possible by measuring alpha-fetoprotein in the amniotic fluid at 14-16 weeks and by prenatal ultrasound examination.

3-**occult spinal dysraphism:**

I. Dermoid tumors.
II. Diastematomyelia.



III. Intraspinal lipoma.

Embryology:

Embriologicaly,the NTD abnormality manifest between 3 and 4 weeks of gestation, at this point of development, the neural plate folds into the neural tube a process called **neurulation**.

Etiology:

Multiples causes can be contributed to the etiology of NTD include:

1. Chromosomal abnormalities.

Down syndrome, klinefelters syndrome.

2. Enviromental factors (teratogens) like:

- Chemical agents
- Drugs like valproic acid.
- Hypothermia
- Hypoxia
- Infections as TORCH.
- Radiation.
- Trauma.
- Diabetes.

DNA synthesis and cell division

3. folic acid deficiency of pregnant mother can attribute to NTD development in fetuses.

Associated anomalies:

NTD can be associated with different poly systemic abnormalities involving different body areas manifested as following:

1-brain:

- Hydrocephalus, 80% of children with meningomylocele develop hydrocephalus at time of presentation or later on.
- Chiari malformation.
- Cerebellar dysgenesis.
- Agenesis of corpus callosum

2-skull anomalies:

- Small posterior fosse.
- Enlarged foramen magnum.
- Low attachment of tentorium.

3-vertebral anomalies:

- Hemi vertebra.
- Butterfly vertebra.

- Vertebral fusion.

4-spinal cord anomalies:

- Chiari malformation.
- Hydromylia.
- Doubling of spinal cord.
- Intraspinal lipoma.

5-systemic anomalies:

- Gastrointestinal tract.
- Genitourinary tract.
- Cardiovascular.
- Rib absence or duplication or bifurcation.

Clinical features of NTD :

1-Hydrocephalus.

2-Cutaneous manifestation:

- Midline dermal patches.
- Subcutaneous lipoma.
- Capillary hemangioma.
- Dermal pit or sinuses.
- Hypertrichosis.

3-sensorimotor deficits:

- Motor weakness either unilateral or bilateral includes atrophy of buttocks, calves, feet.
- Hammer toe.
- Deep tendon reflexes, abscent on affected side.
- Insensitive pin prick sensation may be confined to the perianal area.
- Gait abnormality in older children.

4-sphincteric disturbance, as bowel and bladder dysfunction.

- Bowel: as continuous leakage of patulous anus.
- Bladder: dribbling of small amount of urine which increases with crying or movement. in older children,urgency,frequency, incontinences and frequent UTI.

5-foot anomalies:

- Club foot.

- Varus or valgus deformity of foot.

6-spinal deformity:

- Scoliosis.
- Kyphosis.
- Kyphoscoliosis.

7-trophic changes: of the lower extremities from sympathetic denervation, for example, shiny skin, hair loss, nails changes.

Radiology:

1-plain x-ray of spine:

Show a wide range of vertebral abnormalities including spina bifida, hemivertebrae, abnormal spine curves.

2-MRI:

Give details about NTD, extent and location of lipoma, thickened filum, and spinal cord anomalies.

3-CT scan:

Of brain to examine for hydrocephalus and brain deformities.

4-Skull x-ray:

For skull abnormalities.

5-ultrasound:

Searching for associated anomalies in CVS, GIT, and genitourinary deformities.

6-Electromyogram EMG:

For specific nerve root, spinal nerves and muscular pathology.

7-Echocardiography

For ASD,VSD and other cardiac anomalies.

Treatment:

Surgery followed by physiotherapy and rehabilitation still the corner stone in the treatment.

Surgery in form of exploration of sac and neck of cyst, excision of cyst, reducing intact neural tissues and water tight closure of meninges followed by layered (multi layer facial, subcutaneous and skin) closure.

Treatment also should be directed towards the associated polysystemic deformities accordingly. The patient must be closely monitored and followed clinically and by imaging assessment to detect ant possible complication.

Intussusception

Dr. Hayder N. Hassan
Pediatric surgeon

It is one of the most common causes of intestinal obstruction in infants and toddlers, it occurred when one portion of the bowel enters into immediately adjacent one commonly it is proximal into distal.

Etiology:

intussusception divided according to causes as primary and secondary
The primary (idiopathic) type:
(1) can occur at any age, but mostly between (5-9) months of life. more than half of cases occur in the 1st year of life
(2) occur with the wake of the upper respiratory tract infection and episode of gastroenteritis.
Secondary type: (1): in this case a definite anatomical lead point had been found.
(2): its incidence is 2-12% of cases and usually present after 2 years of age.
(3): the most frequent lead point is Meckel's diverticulum, others include: appendix, polyp, carcinoid, submucosal hemorrhage, foreign body and tumors.

Anatomical classification:

Ileocolic (more than 80%of cases), ileoileal, cecocolic , colocolic and jejunojejunal types.

Clinical presentation: the patient is usually well nourished, healthy infant and 2/3 of cases are male.
1- there is sudden attacks of acute abdominal crump described as screaming attacks during which the patient may stiffen and pull his legs up into the abdomen , with breath holding , the attacks cease suddenly as it started , the patient between the attacks is comfortable and may sleep ,but then become lethargic.

2- vomiting: it become universal in later stages and it become bilious due to intestinal obstruction.
3-normal or small bowel motion appeared initially due to straining and bowel evacuation distal to the site of obstruction. Later, red currant jelly bowel motion appear which is mixture of blood with mucus secretion due to ischemia.

On exam:

1- the vital signs are normal initially, but later tachycardia fever, dehydration even septic shock may occur.
2- abdominal exam shows: intraabdominal mass and empty RIF (dances sign)
3- per rectal (pr) exam. shows red currant jelly, and even may touch the intussusception mass.
4- prolapsing mass though the anus, which is a grave sign and must be differentiated from simple rectal prolapse.

Investigation:

1- plain abdominal radiograph shows: multiple air/fluid levels, soft tissue mass, pnuemoperitonium, abnormal gas distribution.
2- U/S shows target sign (in transverse section) and psuedokidney sign (in longitudinal view).
3- contrast enema (diagnostic and therapeutic) shows claw sign.
4- CT scan: intraluminal mass with layered appearance.

Treatment: 1- resuscitation

2- reduction of intussusception. Which done either by operative or nonoperative method.
A- nonoperative method either by: hydrostatic reduction or by pneumatic reduction.
B- operative method either by laparoscopy or traditional laparotomy.

Infantile hypertrophic pyloric stenosis

Dr. Hayder N. Hassan
Pediatric surgeon

It is the most common surgical cause of vomiting in infancy.

Incidence: its incidence is increasing, male affected 4 times than female , the 1st born male baby and family history are the risk factors.

Etiology: the true etiology is unknown; possible theories includes:

1- congenital redundancy of pyloric mucosa.
2- Abnormalities of local enteric innervation.
3- Prenatal exposure to erythromycin or azythromycin.

Clinical presentations: typically the patient is 2-8 wks of age.
The child has postprandial, forceful, nonbilious vomiting commonly refer to as
Projectile. And the baby is typically hungry after vomiting eager to eat, only to vomit once again.
Diagnosis:
Clinical exam. Shows:

1- Patient generally looks malnourished with or without dehydration'
2- scaphoid abdomen.
3- Gastric peristaltic wave can be seen though the abdominal wall from left to right.
4- Pyloric mass can be palpated in the epigastric region, its palpation as olive confirm the diagnosis and no further investigation needed.

Investigations:

A-Ultrasonography:shows

1- pyloric mass with muscle thickness(>4mm) and pyloric length of (>16mm).
2- Narrow pyloric channel.
3- Redundant thickened mucosa.

B- upper GI contrast study shows:

1- string sign (narrow pyloric channel)
2- shoulder sign (impression of the pyloric into the stomach)

Treatment:
preoperative preparation
Serum electrolytes should always be determined. The most common biochemical abnormalities is hyponatremic hypochloremic hypokalemic metabolic alkalosis with paradoxical acidurea.

How this occurred? - the loss of gastric secretion due to vomiting lead to dehydration, as a result, through aldosterone stimulated absorption, K+ is excreted in the urine in attempt to conserve Na+, and as K+depletion worsens, then Na+ absorption across the renal tubules is achieved in exchange with hydrogen ion (H+) , there by creating paradoxical acidurea.

Initially, a 10-20 ml/kg bolus of normal saline should be given if the electrolytes are abnormal .then, D5/1/2NS with 20-30 mEq/L of kcl is started at a rate of 1.25 to 2 times the calculated maintenance rate.
Electrolytes should be checked every six hours until they normalize and the alkalosis has resolved
Preoperative placement of NG tube is controversy.

Surgical treatment
is done by Ramstedt`s operation since 1912 which is extramucosal muscle-splitting pyloromyotomy operation.

Postoperative complications:

1- Mucosal perforation.
2- persistant vomiting beyond 48hrs mostly due to gastric atony.
3- Duodenal injury
4- Incomplete pyloromyotomy.
5- wound infection and dehiscence 1%

The acute scrotum

It is a sudden onset of painful scrotum that can be accompanied by swelling, erythema and systemic sings of illness.

The term acute scrotum includes a wide spectrum of cases that ranging from those which required immediate surgical intervention to those which required only reassurance.

Several conditions cause a red, swollen and painful scrotum, with wide variations in speed of onset, rate of progression and local signs and the severity of pain. The differential diagnosis is very important as any delay or wrong diagnosis can lead to testicular loss. About 70% of cases not required surgery.

Torsion of the testis

Testicular torsion is not the most common cause of an acute scrotum (16%), but it is the most important. The spermatic cord undergoes torsion, obstructing the spermatic vessels, and is a surgical emergency because of the high incidence of testicular infarction if the cord is not untwisted promptly. The risk of torsion is greatest just after the testis enlarges at puberty in 12–16-year-olds. Also, the risk is increased in neonatal period and in unoperated undescended testes.

Two kinds of torsion occur:

1 Intratunical (or intravaginal), the more common.

2 Extratunical (or extravaginal) torsion is rare.

Clinical signs

The onset is usually sudden, with pain in the testis and/or ipsilateral iliac fossa, nausea and vomiting.

A horizontal lie of the testis when the child stands, often referred to as a bell-clapper testis, indicates the possibility of torsion and should be taken as an indication for exploration and testicular fixation.

The swollen testis and epididymis are severly tender (unless already necrotic) and may be partially obscured by overlying scrotal oedema and an effusion into the tunica (reactive hydrocele).

Treatment

Urgent exploration of the scrotum is arranged to untwist the testis and epididymis.If the testis is completely necrotic, it should be removed.

Torsion of an appendage

Torsion of a testicular appendage is the most common cause of the acute scrotum in prepubertal boys. Testicular (or epididymal) appendages are remnants of the embryonic Mu ́ llerian ducts (that form the uterus and fallopian tubes in females) and are present in about 90% of boys. A small tender lump at the upper pole of the testis is diagnostic.

Clinical signs

The boy complains of severe pain in his scrotum. A blue-black spot (the infarcted appendage with secondary hemorrhage) may be seen through the skin of the scrotum near the upper pole of the testis: palpation of it causes extreme pain, whereas palpation of the testis itself causes minimal discomfort. It may be impossible to distinguish torsion of a testicular appendage from testicular torsion once a secondary hydrocele has developed.

Treatment

If testicular torsion cannot be excluded on clinical examination, urgent exploration is mandatory. At operation, the torted appendix testis is removed, which provides relief of symptoms and prevents recurrence. However, if a torted appendage can be diagnosed on the basis of clinical findings, non-operative treatment with effective analgesia is a valid alternative.

Epididymitis or epididymo-orchitis

Epididymo-orchitis is rare in childhood and virtually never occurs between 6 months of age and puberty. Although it is common practice to refer to inflammatory conditions in the scrotum as epididymo-orchitis, the inflammation is usually confined to the epididymis. The most common causative bacterium in children is Escherichia coli, carried by retrograde flow along a patent vas deferens from the urinary tract. Predisposing factors for bacterial infection include abnormalities of the urinary tract and urethral instrumentation.

Clinical signs

The usual findings are those of an acute scrotum in a baby or adolescent. A lax secondary hydrocele is common, and bilateral signs are particularly suggestive of epididymitis. Examination of the urine may show pyobacteriuria.

Young children with epididymitis due to urinary organisms should have a renal ultrasound scan after the epididymitis has subsided, and some may require also a micturating cystourethrogram. These investigations aim to identify anomalies of the lower urinary tract before irreversible damage to the kidneys has occurred.

Differential diagnosis

True acute orchitis is very uncommon, but may occur in mumps, Henoch–Schönlein purpura (HSP) or septicaemia. Mumps orchitis is extremely rare prior to puberty. Malignant infiltration of the testis is rare also, but does occur occasionally in leukaemia or with a primary neoplasm (embryonic adenocarcinoma, seminoma or a benign tumour of the Leydig cells).

Treatment

Treatment of epididymitis consists of rest, antibiotics (e.g. co-trimoxazole, nitrofurantoin), a high fluid intake.

Idiopathic scrotal oedema

In this condition, there is rapidly developing scrotal oedema, which may then spread to the inguinal region, penis and foreskin and/or the perineum. The scrotum is symmetrically swollen, pale pink or red, and there is slight discomfort rather than acute pain. The pathology involves the skin and may represent allergic inflammation.

Careful palpation reveals non-tender testes that are normal in size and position. The oedema subsides in 1–2 days, but may occasionally recur some weeks later. There may be a history of allergy or of playing outside at the onset; a bite from an insect or a spider is a probable cause in some, but as a rule, the history is inconclusive.

Inguinal lymphadenitis

The superficial inguinal lymph nodes drain the lower limbs, the perineum, the buttocks and the perianal region – all common sites of minor skin infections in the nappy area in infants. Infections often reach the inguinal nodes, which become enlarged and may form an abscess after the initial focus has disappeared.

Anorectal malformation (ARM)

Ass. Proff. Hayder N. Hassan
consultant Pediatric surgeon

It is congenital malformation that affect the more distal part of the GI tract.

Incidence & Terminology:

- ARM occur in ONE of every 4000-5000 newborns.
- Slightly more common in males.
- Most frequent defect in male is rectourethral fistula while in female is rectovestibular fistula.
- Rectovaginal fistula is a rare defect , present in < 1% of all cases .

Classification:

Anorectal malformations represent a wide spectrum of defects. The terms “low,” “intermediate,” and “high” are arbitrary and not useful in therapeutic or prognostic terms.

Male defects	female defects
Recto perineal fistula	Recto perineal fistula
Recto urethral bulbar fistula	Recto vestibular fistula
Recto urethral prostatic fistula	Cloaca
Recto bladder neck fistula	Complex malformations
Imperforated anus without fistula	Imperforated anus without fistula
Rectal atresia	Rectal atresia

We`ll discuss the some common types in male and female.

Male defects:

Recto perianal fistula : which is previously called low type in such a case:

1- the rectum located within the sphincter but its lowest part is anteriorly mislocated.
2- Diagnosis is clinical by inspection, there is a cutaneous fistula.
3- Treatment by anoplasty without protective colostomy in the 1st 48 hrs of life.

Rectourethral fistula:

1- it is the most frequent defect in the male patients .

2- the fistula may be located at the lower (bulbar) or the higher (prostatic) part of urethra.

The bulbar type associated with good functional prognosis because there is good muscle quality, well developed sacrum, and prominent midline groove, while the prostatic type associated with poor functional prognosis.

If the infant passes meconium through the urethra, this is an unequivocal sign of a rectourinary fistula

diagnosis by

a- history: there is no passage of meconium, no anal opening and may be meconium coming with urine .

b- on exam. Inspection shows there is no anus and meconium may be seen coming with urine.

c- Investigation: during the 1st 24hrs insert NG tube and send for abdominal U/S to search for any associated anomalies .after the 1st 24 hrs. send for cross –table lateral plain x-ray with the patient in prone position where it shows the distal extension of air within the rectum whether it is beyond or above the coccyx.

treatment:by

a- resuscitation by:

1- rest of the bowel (NG tube and no thing by mouth)

2- correction of fluid and electrolytes disturbances.

3- IV antibiotics
4- Vit. K if there is signs of bleeding tendency.

b- decompression by doing divided descending colostomy, then discharge the patient home with stoma.

c- After 4 weeks and the patient has a good weight do definite procedure which is (PSARP)posterior sagittal anorectoplasty.

Rectobladderneck fistula

- The rectum opens into the bladderneck.
- Poor Prognosis for bowel control/ poorly developed muscles
- Deformed and short sacrum
- Flat perineum

Female defects:

A- Rectoperineal fistula in females

Equivalent to the defect in males.

B- **rectovestibular fistula** :

1- it is the most common female defect with excellent functional prognosis.
2- Diagnosis is clinical one by inspection,by finding a third hole in the vestibule
3- Treatment by doing protective colostomy then definite procedure.

C- **Cloaca**: It represents the extreme in the spectrum of complexity of female malformations.

it is a defect in which the rectum, vagina, and urinary tract meet and fuse, creatinga single common perineal channel, the diagnosis is clinical by inspection.

- Associated Defects:
- Sacrum and Spine.
- Tethered Cord
- VACTERL (vertebral, anal, cardiac,tracheoesophageal renal and limb).
- Associated GU defects varies from 20-54%.
- The higher the malformation, the more frequent are the associated urologic anomalies.

Newborn Management

- Thorough perineal inspection
- No decision whether colostomy or primary repair before 24 hours of life.
- I.V fluid, AB, NG decompression and evaluation for associated defects (cardiac ECHO, Renal and spinal US
- If there is signs of rectoperineal fistula ➔ Anoplasty.
- No meconium in the perineum, we recommend obtain cross-table lateral radiograph in prone position.
- If air in the rectum is seen distal to the coccyx, consider PSARP without protective colostomy.
  If the rectal gas does not extend beyond the coccyx or the patient pass meconium in urine, do colostomy

Complications after PSARP

- Wound infection
- Wound separation
- Anal strictures
- Constipation
- Rectal prolapse
- Transient femoral nerve palsy.
- Urologic injuries

CONGENITAL DIAPHRAGMATIC HERNIA

Ass.proff. Hayder N. Hassan
Consultant pediatric surgeon

The diaphragm develops from four embryonic structures:
1 The septum transversum
2 The left and right pleuro-peritoneal membranes
3 Dorsal oesophageal mesentery
4 Somites at cervical segments 3–5

Congenital diaphragmatic hernia results from failure of formation or fusion of the components of the diaphragm, such that abdominal contents may move through a defect into the thoracic cavity. Sometimes failure of muscularisation may produce a thin, weak diaphragm, referred to as an eventration of the diaphragm.

Types:

The **Bochdalek type** is the most common(85%) variety of congenital diaphragmatic hernia (1 in 5000 live births) and results from a defect in the postero-lateral aspect of the diaphragm. During intra-uterine development, the small bowel, stomach, spleen and left lobe of the liver may pass through the defect in the diaphragm into the chest. Lung development is also abnormal in fetuses with congenital diaphragmatic hernia, with hypoplastic lungs and pulmonary vasculature.

The **Morgagni** (retrosternal) type of diaphragmatic hernia is rare (13%) and results from a defect in the anterior midline, just behind the sternum. It usually contains part of the colon or small bowel and, less commonly, part of the liver.
bilateral (2%).

Oesophageal hiatal herniae may also occur and usually produce symptoms of gastro-oesophageal reflux.

Clinical features

Antenatal diagnosis

The widespread use of obstetric sonography has led to an increase in the frequency of antenatal diagnosis of CDH, which is established by demonstration of the abdominal viscera in the chest. Three easily detectable features – polyhydramnios, mediastinal shift and the absence of an intra-abdominal stomach bubble – should prompt a more careful search for herniated abdominal organs in the chest. Polyhydramnios is present in about 80% of the pregnancies with fetuses who have CDH and has also been associated with poor outcome.

Postnatal diagnosis

Postnatally, the most severely affected babies present with respiratory distress (cyanosis, tachypnea and sternal recession) at birth. Other neonates develop cyanosis, tachypnoea and grunting respirations with increased respiratory effort, within minutes or hours after birth. Physical examination reveals a scaphoid abdomen, an increased anteroposterior diameter of the thorax and mediastinal shift. Breath sounds are absent on the affected side and heart sounds are on the right side.
Associated congenital anomalies may also be seen or revealed on further examination.
It occurs in up to 40%, but most are minor and do not affect survival, for example, undescended testes. The most common serious abnormalities are heart defects, which affect 20–25% of patients.

Investigation

Diagnosis of CDH is made postnatally by plain radiography of the chest and abdomen by demonstration of air-filled loops of the bowel in the chest and a paucity of gas in the abdomen The heart is deviated to the right. The diaphragmatic margin is absent, there is a mediastinal shift to the opposite side and only a small portion of the lung may be seen on the ipsilateral side.
Sometimes the appearance may be difficult to distinguish from basal lung cysts, in which case a repeat chest x-ray is performed after a nasogastric tube has been inserted, the tip of which may be seen in the chest. Alternatively, a barium study will show bowel within the thoracic cavity when there is a diaphragmatic hernia.

Treatment

Where antenatal ultrasonography has identified a diaphragmatic hernia, the best outcomes are achieved if the neonate is transferred to a tertiary pediatric surgical center prior to birth. This is because these neonates may develop severe pulmonary distress very quickly after birth, making subsequent transfer difficult and potentially dangerous.

Initial treatment involves intensive cardiorespiratory support and insertion of a nasogastric tube to prevent bowel dilatation within the chest. Care must be taken to avoid hyperinflation and barotrauma of the small, hypoplastic lungs. Ventilation with a facemask (‘bagging’) should be avoided as this may force air into the stomach, increasing its volume at the expense of the already compromised lungs. Vigorous endotracheal ventilation should also be avoided because of the risk of causing barotrauma or tension pneumothorax, which may lead to the rapid demise of the neonate. Exogenous surfactant provides no specific benefit in newborns with diaphragmatic hernia. The key to success is careful gentle ventilation that minimises injury to the hypoplastic lungs.
Sudden deterioration of the neonate’s condition during initial resuscitation or during transport suggests the development of a tension pneumothorax, and this may necessitate prompt release by needle aspiration or insertion of an intercostal drain. The primary focus of management of the neonate with congenital diaphragmatic hernia is achieving adequate ventilation with control of pulmonary hypertension. To reduce the risk of injury to hypoplastic lungs in these patients, special ventilatory strategies may be required such as high-frequency oscillation or in very selected cases extracorporeal membrane oxygenation.

Surgery to return the herniated contents to the abdominal cavity and to repair the defect in the diaphragm is only performed once the neonate’s condition has stabilized. Surgery is typically deferred for at least 24 h after birth, but stabilization may not be achieved for many days. Surgery generally involves repair through a transverse or subcostal abdominal incision on the same side as the defect. Thoracoscopic surgery has a role in selected cases,

Survival rates post-surgery of more than 80% are now being achieved, and may be higher in specialist centers. The major cause of death remains pulmonary hypoplasia and pulmonary hypertension

Key points:

• Congenital diaphragmatic hernia is diagnosed on antenatal ultrasonography or by chest x-ray in a newborn with a barrel chest, scaphoid abdomen and/or respiratory distress.
• Ventilatory support, especially during transport, should be the minimum required to prevent deterioration, as hyperinflation with secondary barotrauma is a significant complication.
• Sudden deterioration is usually the result of a tension pneumothorax

CONGENITAL EVENTRATION OF THE DIAPHRAGM

Eventration of the diaphragm has been described as an abnormally high or deviated position of all or part of the hemidiaphragm. Eventration may be congenital or acquired as a result of phrenic nerve palsy. Congenital eventration is a developmental abnormality which results in muscular aplasia of the diaphragm. In acquired eventration, the diaphragm, which initially had fully developed musculature, becomes atrophic secondary to phrenic nerve damage and disuse.

Clinical features range from being asymptomatic to severe respiratory distress. Patients may present later in infancy with repeated attacks of pneumonia, bronchitis or bronchiectasis. Occasionally, patients present later in childhood with gastrointestinal symptoms of vomiting or epigastric discomfort. In patients with phrenic nerve palsy, there may be a history of difficult delivery. They may present with tachypnoea, respiratory distress or cyanosis.

Physical examination reveals decreased breath sounds on the affected side, mediastinal shift during inspiration and a scaphoid abdomen. The diagnosis of eventration is usually made on a chest X-ray. Frontal and lateral chest X-rays will show an elevated diaphragm with a smooth,unbroken outline.
Fluoroscopy is a useful investigation for differentiating a complete eventration from a hernia. Paradoxical movement of the diaphragm is seen if complete eventration is present.
Ultrasonography is the most useful study in the diagnosis of eventration of the diaphragm and for identification of abdominal organs underneath the eventration., radioisotope scanning and computed tomography scans but these are rarely required.

Symptomatic patients, especially those with respiratory distress, need prompt supportive care with endotracheal intubation and ventilation with humidified oxygen to minimize the diaphragmatic excursions. A nasogastric tube is passed to decompress the stomach and intravenous fluids are commenced. Surgery is undertaken once the patient’s condition is stabilized.

Se: 1001/2
Im: 1001/1
CHEST